

DEX-0117

DEX-116

AD



9-721183

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/12, C07K 14/47, 16/18, C12N 15/63, 5/10, 1/21, C12Q 1/68		A1	(11) International Publication Number: WO 98/33915 (43) International Publication Date: 6 August 1998 (06.08.98)
(21) International Application Number: PCT/US98/01804 (22) International Filing Date: 3 February 1998 (03.02.98) (30) Priority Data: 60/037,080 3 February 1997 (03.02.97) US (71) Applicant (for all designated States except US): HUMAN GENOME SCIENCES, INC. [US/US]; 9410 Key West Avenue, Rockville, MD 20850 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): JI, Hongjun [CN/US]; 13144 Country Ridge Drive, Germantown, MD 20874 (US). ROSEN, Craig, A. [US/US]; 22400 Rolling Hill Road, Laytonsville, MD 20882 (US). (74) Agents: STEFFE, Eric, K.; Sterne, Kessler, Goldstein & Fox P.L.L.C., Suite 600, 1100 New York Avenue, N.W., Washington, DC 20005-3934 (US) et al.		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments. With an indication in relation to a deposited microorganism furnished under Rule 13bis separately from the description. Date of receipt by the International Bureau: 10 March 1998 (10.03.1998)	

(54) Title: BREAST CANCER SPECIFIC GENE 1

(57) Abstract

The present invention relates to a BCSG1 protein. In particular, isolated nucleic acid molecules are provided encoding the human BCSG1 protein. BCSG1 polypeptides are also provided as are vectors, host cells and recombinant methods for producing the same. Also provided are diagnostic methods for detecting breast cancer.

```
10          30          50
CACGAGCCACCATGGATGTTTTCAGAGAGGCGCTTCCATGCCAAGAGGCGCOTGGTGG
M D V F K K Q P S I A K K G V V G

70          90          110
GTGCGCGTGGAAAGAGCCAGCAGCGGCGTACGAGAGCACTGAGAGAGCAAGAGAGG
A V E K T K Q Q V T E A A E K T K E G V

130          150          170
TCATGTATGTGGAGCCAGACCAAGGAGAAATGTTATACAGAGCGGTGACCTCACTGGCCG
M T V G A K T K E H V V Q S V T S V A E

190          210          230
AGAAGACCAAGGAGCAAGCCACCGCGTGAAGCAAGCGCTGTGTGAGCAGCGTCAACACTG
K T K E Q A N A V S K A V V E S V H T V

250          270          290
TGCCACCAAGACCGTGGAGGAGGCGGAGAACATCCCGGTACCTCCCGGCGTGTGGCCA
A T K T V E E A E N I A V T S G V V R K

310          330          350
AGGAGGACTTGAAGCCATCTCCCGCCCAACAGAGGCGTGAAGCATCCAAGAGAGAGAGG
E D L R P S A P Q Q E G E A S K E K E E

370          390          410
AAGTGGCAGAGGAGGCCAGAGTGGGAGGAGCTAGAGGGCTACAGGCGCAGCGTGGATGAC
V A E E A Q S G G D *

430          450          470
CTGAAGAGCGCTCTCTCTGCTTGGACACCATCCCTCTAGCACAAGAGTCCCGCGCTT
GAGTGACATGCGGTCGCCACGCTCTGCGCTGCTCTGCGTGGACACCTTGGCGTTGCTC
550
ACCTGTGCTG
```

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Breast Cancer Specific Gene 1

Field of the Invention

The present invention relates to a novel breast cancer specific marker. More specifically, isolated nucleic acid molecules are provided encoding a human breast cancer specific gene 1 (BCSG1). BCSG1 polypeptides are also provided, as are vectors, host cells and recombinant methods for producing the same. Also provided are diagnostic methods for detecting breast cancer. The invention further provides an isolated BCSG1 polypeptide having an amino acid sequence encoded by a polynucleotide described herein.

Related Art

More than 190,000 new cases of breast cancer are diagnosed in the United States every year, with incidence increasing by approximately 1% annually (Goldhirsch, A., *JNCI* 97:1141-1145 (1995); Emster, V.L., *et al.*, *JAMA* 275:913-918 (1996)). Studies linked to the discovery of new genetic markers and additional risk factors could provide new information that fits into the complex patient management issues surrounding breast cancer. Many new prognostic and predictive factors have been proposed and studied for breast cancer. HER 2/neu positive tumors respond poorly to endocrine treatment (Allred D.C., *et al.*, *J. Clin Oncol.* 10:599-605 (1992); Gusterson B.A., *et al.*, *J. Clin Oncol.* 10:1049-56 (1992)). p53 alteration has an indication of poorer prognosis and poor response to tamoxifen (Bergh J., *et al.*, *Nature Medicine* 10:1029-34 (1995); Elledge R.M., *et al.*, *Breast Cancer Res Treat* 27:95-102 (1993)). The lack of Nm23 expression has an indicative value of metastatic potential and poor prognosis in invasive ductal carcinoma (Steeg P.S., *et al.*, *Breast Cancer Res Treat* 25:175-87 (1993)). Cathepsin D, a protease suggested to have a role in breast cancer, appears to affect the potential for invasive growth (Velculescu, V.E., *et al.*, *Science* 270:484-7 (1995); Schena, M., *et al.*, *Science* 270:467-70 (1995); M.L. Angerer & R.C.

Angerer, In: *In situ hybridization*, D. Rickwood and B.D. Hames (ed.). London: LRL Press., (1992), pp. 15-32; Ferno M., *et al.*, *Eur J. Cancer* 30A:2042-8 (1994)). Positive immunostaining of tumor sections with Factor VIII antibodies seems to be a marker for angiogenesis (Klijn J.G.M., *et al.*, *Breast Cancer* 18:165-98 (1993); Harris A.L., *et al.*, *Eur J. Cancer* 31A:831-2 (1995); Gasparini G., *et al.*, *JNCI* 85:1206-19 (1993) (errata *JNCI* 85:1605 (1993))). It has been postulated that these tumors are targets for anti-angiogenesis drug treatment. Expression of the *mdr-1* gene is proposed to be an indicator of multidrug resistance (Harris A.L., *et al.*, *Eur J. Cancer* 31A:831-2 (1995); Gasparini G., *et al.*, *JNCI* 85:1206-19 (1993) (errata *JNCI* 85:1605 (1993))). Poor response to endocrine therapy has been indicated for uPA/PAI-1, a plasminogen activator/inhibitor (Foekens J.A., *et al.*, *JNCI* 87:751-6 (1995)). Also receiving major attention are the familial breast cancer related genes, BRCA1 and BRCA2 (Miki, Y., *et al.*, *Science* 266:66-71 (1994); Wooster, R., *et al.*, *Science* 265:2088-2090 (1994); Futreal, P.A., *et al.*, *Science* 266:120-122 (1994)).

Thus, the onset and progression of breast cancer is accompanied by multiple genetic changes that result in qualitative and quantitative alterations in individual gene expression (Porter-Jordan, K. & Lippman, M.E., *Hematol. Oncol. Clin. N. Am.* 8:73-100 (1994)). Many of these quantitative genetic changes may manifest themselves as alterations in the cellular complement of novel transcribed mRNAs. Identification of these mRNAs could provide clinically useful information for patient management and prognosis while enhancing our understanding of breast cancer pathogenesis.

Identification of quantitative changes in gene expression that occur in the malignant mammary gland may yield novel molecular markers which may be useful in the diagnosis and treatment of human breast cancer. Several differential cloning methods, such as differential display polymerase chain reaction and subtractive hybridization, have been used to identify the genes differentially expressed in breast cancer biopsies, as compared to normal breast tissue controls (Watson, M.A. & Fleming, T.P., *Cancer Res.* 54:4598-4602 (1994); Sager, R., *et*

al., *FASEB J.* 7:964-970 (1993); Chen, Z. & Sager, R., *Mol. Med.* 1:153-160 (1995); Zhang, M., *et al.*, *Cancer Res.* 55:2537-2541 (1995); Zou, Z., *et al.*, *Science* 263:526-529)). However, these investigations have involved the relatively time- and labor-intensive steps of subcloning, library screening, and cDNA sequencing of individual genes (Sager, R., *et al.*, *FASEB J.* 7:964-970 (1993); Liang, P., *et al.*, *Cancer Res.* 52:6966-6968 (1992)).

Although pathological endpoints such as tumor size, lymph node status and status of estrogen receptor and progesterone receptor remain the most useful guides in prognosis and selecting treatment strategies for breast cancer (Manning, D.L., *et al.*, *Acta Oncol.* 34:641-646 (1995)), there is still a need to further investigate the molecular mechanisms that determine the properties of an individual tumor e.g., probability of metastasis. While numerous prognostic factors have been identified, few have contributed to defining clinical response to therapy.

Summary of the Invention

The present invention provides isolated nucleic acid molecules comprising a polynucleotide encoding the BCSG1 polypeptide having the amino acid sequence shown in Figure 1 (SEQ ID NO:2) or the amino acid sequence encoded by the cDNA clones deposited in a bacterial host as ATCC Deposit Number 97175 on June 2, 1995 or as ATCC Deposit Number 97856 on January 23, 1997.

The present invention also relates to recombinant vectors, which include the isolated nucleic acid molecules of the present invention, and to host cells containing the recombinant vectors, as well as to methods of making such vectors and host cells and for using them for production of BCSG1 polypeptides or peptides by recombinant techniques.

In accordance with another aspect of the present invention, there is provided a method of and products for diagnosing breast cancer metastases by detecting an altered level of a polypeptide corresponding to the breast specific

genes of the present invention in a sample derived from a host, whereby an elevated level of the polypeptide indicates a breast cancer diagnosis.

The present invention further relates to antibodies specific to the polypeptides of the present invention, which may be employed to detect breast cancer cells or breast cancer metastasis.

The polynucleotides and polypeptides described herein are useful as markers for breast cancer.

Brief Description of the Figures

Figure 1 shows the nucleotide (SEQ ID NO:1) and deduced amino acid (SEQ ID NO:2) sequences of BCSG1. The protein has a deduced molecular weight of about 14.2 kDa. The predicted amino acid sequence of the BCSG1 protein is also shown.

Figure 2 shows the differential cDNA sequencing approach. Messenger RNAs from normal and diseased tissues were extracted and used for making the cDNA libraries. These libraries are searched by EST method involving automated DNA sequence analysis of randomly selected cDNA clones. The ESTs with overlapping sequences were grouped into unique EST groups. Each unique EST group, which does not overlap to each other in sequence, was analyzed for its relative expression by examining the number of expressed individual EST in the libraries of normal vs diseased tissues. Three EST groups are listed. Blue EST group represents gene that is equally expressed in both libraries. Green EST group represents gene that is more expressed in normal library compared to diseased library. Red EST group represent gene that is more expressed in diseased library compared to normal library.

Figure 3 shows a schematic representation of the pHE4-5 expression vector (SEQ ID NO:10) and the subcloned BSCG-1 cDNA coding sequence. The locations of the kanamycin resistance marker gene, the BSCG-1 coding sequence, the oriC sequence, and the *lacIq* coding sequence are indicated.

Figure 4 shows the nucleotide sequence of the regulatory elements of the pHE promoter (SEQ ID NO:11). The two *lac* operator sequences, the Shine-Delgarno sequence (S/D), and the terminal *Hind*III and *Nde*I restriction sites (*italicized*) are indicated.

5

Detailed Description

The present invention provides isolated nucleic acid molecules comprising a polynucleotide encoding a BCSG1 polypeptide having the amino acid sequence shown in Figure 1 (SEQ ID NO:2), which was determined by sequencing a cloned cDNA. The BCSG1 protein of the present invention shares sequence homology
10 with human AD amyloid. The nucleotide sequence shown in Figure 1 (SEQ ID NO:1) was obtained by sequencing the 184,497 clone, which was deposited on January 23, 1997 at the American Type Culture Collection, 12301 Parklawn Drive, Rockville, Maryland 20852, and given accession number 97856. The deposited clone is contained in the pBluescript SK(-) plasmid (Stratagene, La
15 Jolla, CA). The BSCG-1 gene was also deposited on June 2, 1995 at the American Type Culture Collection, 12301 Parklawn Drive, Rockville, Maryland 20852, and given accession number 97175.

Nucleic Acid Molecules

Unless otherwise indicated, all nucleotide sequences determined by
20 sequencing a DNA molecule herein were determined using an automated DNA sequencer (such as the Model 373 from Applied Biosystems, Inc.), and all amino acid sequences of polypeptides encoded by DNA molecules determined herein were predicted by translation of a DNA sequence determined as above. Therefore, as is known in the art for any DNA sequence determined by this
25 automated approach, any nucleotide sequence determined herein may contain some errors. Nucleotide sequences determined by automation are typically at least about 90% identical, more typically at least about 95% to at least about

99.9% identical to the actual nucleotide sequence of the sequenced DNA molecule. The actual sequence can be more precisely determined by other approaches including manual DNA sequencing methods well known in the art. As is also known in the art, a single insertion or deletion in a determined nucleotide sequence compared to the actual sequence will cause a frame shift in translation of the nucleotide sequence such that the predicted amino acid sequence encoded by a determined nucleotide sequence will be completely different from the amino acid sequence actually encoded by the sequenced DNA molecule, beginning at the point of such an insertion or deletion.

Using the information provided herein, such as the nucleotide sequence in Figure 1, a nucleic acid molecule of the present invention encoding a BCSG1 polypeptide may be obtained using standard cloning and screening procedures, such as those for cloning cDNAs using mRNA as starting material. Illustrative of the invention, the nucleic acid molecule described in Figure 1 (SEQ ID NO:1) was discovered in a cDNA library derived from breast cancer. The gene was also identified in cDNA libraries from brain tissue. The determined nucleotide sequence of the BCSG1 cDNA of Figure 1 (SEQ ID NO:1) contains an open reading frame encoding a protein of 127 amino acid residues, with an initiation codon at positions 12-14 of the nucleotide sequence in Figure 1 (SEQ ID NO:1), and a deduced molecular weight of about 14.2 kDa. The BCSG1 protein shown in Figure 1 (SEQ ID NO:2) is about 54 % identical to non-A β fragment of human Alzheimer's disease (AD) amyloid protein.

As one of ordinary skill would appreciate, due to the possibilities of sequencing errors, the predicted BCSG1 polypeptide encoded by the deposited cDNA comprises about 127 amino acids, but may be anywhere in the range of 110-140 amino acids.

As indicated, nucleic acid molecules of the present invention may be in the form of RNA, such as mRNA, or in the form of DNA, including, for instance, cDNA and genomic DNA obtained by cloning or produced synthetically. The DNA may be double-stranded or single-stranded. Single-stranded DNA or RNA

-7-

may be the coding strand, also known as the sense strand, or it may be the non-coding strand, also referred to as the anti-sense strand.

By "isolated" nucleic acid molecule(s) is intended a nucleic acid molecule, DNA or RNA, which has been removed from its native environment. For example, recombinant DNA molecules contained in a vector are considered isolated for the purposes of the present invention. Further examples of isolated DNA molecules include recombinant DNA molecules maintained in heterologous host cells or purified (partially or substantially) DNA molecules in solution. Isolated RNA molecules include *in vivo* or *in vitro* RNA transcripts of the DNA molecules of the present invention. Isolated nucleic acid molecules according to the present invention further include such molecules produced synthetically.

Isolated nucleic acid molecules of the present invention include DNA molecules comprising an open reading frame (ORF) shown in Figure 1 (SEQ ID NO:1) and DNA molecules which comprise a sequence substantially different from those described above but which, due to the degeneracy of the genetic code, still encode the BCSG1 protein. Of course, the genetic code is well known in the art. Thus, it would be routine for one skilled in the art to generate such degenerate variants.

In another aspect, the invention provides isolated nucleic acid molecules encoding the BCSG1 polypeptide having an amino acid sequence encoded by the cDNA clone contained in the plasmid deposited as ATCC Deposit No. 97856 on January 23, 1997 or contained in the plasmid deposited as ATCC Deposit No. 97175 on June 2, 1995. The invention further provides an isolated nucleic acid molecule having the nucleotide sequence shown in Figure 1 (SEQ ID NO:1) or the nucleotide sequence of the BCSG1 cDNA contained in the above-described deposited clone, the full-length BCSG1 polypeptide lacking the N-terminal methionine or a nucleic acid molecule having a sequence complementary to one of the above sequences. Such isolated molecules, particularly DNA molecules, are useful as probes for gene mapping, by *in situ* hybridization with

chromosomes, and for detecting expression of the BCSG1 gene in human tissue, for instance, by Northern blot analysis.

The present invention is further directed to fragments of the isolated nucleic acid molecules described herein. By a fragment of an isolated nucleic acid molecule having the nucleotide sequence of the deposited cDNA or the nucleotide sequence shown in Figure 1 (SEQ ID NO:1) is intended fragments at least about 15 nt, and more preferably at least about 20 nt, still more preferably at least about 30 nt, and even more preferably, at least about 40 nt in length which are useful as diagnostic probes and primers as discussed herein. Of course, larger fragments 50, 75, 100, 125, 150, 175, 200, 225, 250, 275, 300, 325, 350, 375, 400, 425, 450, 475, 500, 525, 550 nt in length are also useful according to the present invention as are fragments corresponding to most, if not all, of the nucleotide sequence of the deposited cDNA or as shown in Figure 1 (SEQ ID NO:1). By a fragment at least 20 nt in length, for example, is intended fragments which include 20 or more contiguous bases from the nucleotide sequence of the deposited cDNA or the nucleotide sequence as shown in Figure 1 (SEQ ID NO:1).

Preferred nucleic acid fragments of the present invention include nucleic acid molecules encoding epitope-bearing portions of the BCSG1 protein. In particular, such nucleic acid fragments of the present invention include nucleic acid molecules encoding: a polypeptide comprising amino acid residues from about 94 to about 107 in Figure 1 (SEQ ID NO:2); a polypeptide comprising amino acid residues from about 120 to about 127 in Figure 1 (SEQ ID NO:2). The inventors have determined that the above polypeptide fragments are antigenic regions of the BCSG1 protein. Methods for determining other such epitope-bearing portions of the BCSG1 protein are described in detail below.

In another aspect, the invention provides an isolated nucleic acid molecule comprising a polynucleotide which hybridizes under stringent hybridization conditions to a portion of the polynucleotide in a nucleic acid molecule of the invention described above, for instance, the cDNA clones contained in ATCC

Deposits 97856 or 97175. By "stringent hybridization conditions" is intended overnight incubation at 42°C in a solution comprising: 50% formamide, 5x SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 g/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1x SSC at about 65°C.

By a polynucleotide which hybridizes to a "portion" of a polynucleotide is intended a polynucleotide (either DNA or RNA) hybridizing to at least about 15 nucleotides (nt), and more preferably at least about 20 nt, still more preferably at least about 30 nt, and even more preferably about 30-70 nt of the reference polynucleotide. These are useful as diagnostic probes and primers as discussed above and in more detail below.

By a portion of a polynucleotide of "at least 20 nt in length," for example, is intended 20 or more contiguous nucleotides from the nucleotide sequence of the reference polynucleotide (e.g., the deposited cDNA or the nucleotide sequence as shown in Figure 1 (SEQ ID NO:1)). Of course, a polynucleotide which hybridizes only to a poly A sequence (such as the 3' terminal poly(A) tract of the BCSG1 cDNA shown in Figure 1 (SEQ ID NO:1)), or to a complementary stretch of T (or U) residues, would not be included in a polynucleotide of the invention used to hybridize to a portion of a nucleic acid of the invention, since such a polynucleotide would hybridize to any nucleic acid molecule containing a poly (A) stretch or the complement thereof (e.g., practically any double-stranded cDNA clone).

As indicated, nucleic acid molecules of the present invention which encode a BCSG1 polypeptide may include those encoding the amino acid sequence of the polypeptide, by itself; the coding sequence for the polypeptide and additional sequences, such as those encoding an amino acid leader or secretory sequence, such as a pre-, or pro- or prepro- protein sequence; the coding sequence of the polypeptide, with or without the aforementioned additional coding sequences, together with additional, non-coding sequences, including for example, but not limited to introns and non-coding 5' and 3' sequences, such as

the transcribed, non-translated sequences that play a role in transcription, mRNA processing, including splicing and polyadenylation signals, for example - ribosome binding and stability of mRNA; an additional coding sequence which codes for additional amino acids, such as those which provide additional functionalities. Thus, the sequence encoding the polypeptide may be fused to a marker sequence, such as a sequence encoding a peptide which facilitates purification of the fused polypeptide. In certain preferred embodiments of this aspect of the invention, the marker amino acid sequence is a hexa-histidine peptide, such as the tag provided in a pQE vector (Qiagen, Inc.), among others, many of which are commercially available. As described in Gentz *et al.*, *Proc. Natl. Acad. Sci. USA* 86:821-824 (1989), for instance, hexa-histidine provides for convenient purification of the fusion protein. The "HA" tag is another peptide useful for purification which corresponds to an epitope derived from the influenza hemagglutinin protein, which has been described by Wilson *et al.*, *Cell* 37: 767 (1984). As discussed below, other such fusion proteins include the BCSG1 fused to Fc at the N- or C-terminus.

The present invention further relates to variants of the nucleic acid molecules of the present invention, which encode portions, analogs or derivatives of the BCSG1 protein. Variants may occur naturally, such as a natural allelic variant. By an "allelic variant" is intended one of several alternate forms of a gene occupying a given locus on a chromosome of an organism. *Genes II*, Lewin, B., ed., John Wiley & Sons, New York (1985). Non-naturally occurring variants may be produced using art-known mutagenesis techniques.

Such variants include those produced by nucleotide substitutions, deletions or additions, which may involve one or more nucleotides. The variants may be altered in coding regions, non-coding regions, or both. Alterations in the coding regions may produce conservative or non-conservative amino acid substitutions, deletions or additions. Especially preferred among these are silent substitutions, additions and deletions, which do not alter the properties and

activities of the BCSG1 protein or portions thereof. Also especially preferred in this regard are conservative substitutions.

Further embodiments of the invention include isolated nucleic acid molecules comprising a polynucleotide having a nucleotide sequence at least 90% identical, and more preferably at least 95%, 96%, 97%, 98% or 99% identical to (a) a nucleotide sequence encoding the BCSG1 polypeptide having the amino acid sequence in Figure 1 (SEQ ID NO:2); (b) a nucleotide sequence encoding the polypeptide having the amino acid sequence in SEQ ID NO:2, but lacking the N-terminal methionine; (c) a nucleotide sequence encoding the BCSG1 polypeptide having the amino acid sequence encoded by the cDNA clones contained in ATCC Deposit Nos. 97856 or 97175; or (d) a nucleotide sequence complementary to any of the nucleotide sequences in (a), (b) or (c).

By a polynucleotide having a nucleotide sequence at least, for example, 95% "identical" to a reference nucleotide sequence encoding a BCSG1 polypeptide is intended that the nucleotide sequence of the polynucleotide is identical to the reference sequence except that the polynucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence encoding the BCSG1 polypeptide. In other words, to obtain a polynucleotide having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. These mutations of the reference sequence may occur at the 5' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence.

As a practical matter, whether any particular nucleic acid molecule is at least 90%, 95%, 96%, 97%, 98% or 99% identical to, for instance, the nucleotide sequence shown in Figure 1 (SEQ ID NO:1) or to the nucleotides sequence of the

deposited cDNA clone can be determined conventionally using known computer programs such as the Bestfit program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, 575 Science Drive, Madison, WI 53711. Bestfit uses the local homology algorithm of Smith and Waterman, *Advances in Applied Mathematics* 2: 482-489 (1981), to find the best segment of homology between two sequences. When using Bestfit or any other sequence alignment program to determine whether a particular sequence is, for instance, 95% identical to a reference sequence according to the present invention, the parameters are set, of course, such that the percentage of identity is calculated over the full length of the reference nucleotide sequence and that gaps in homology of up to 5% of the total number of nucleotides in the reference sequence are allowed.

The present application is directed to nucleic acid molecules at least 90%, 95%, 96%, 97%, 98% or 99% identical to the nucleic acid sequence shown in Figure 1 (SEQ ID NO:1) or to the nucleic acid sequence of the deposited cDNA, irrespective of whether they encode a polypeptide having BCSG1 activity. This is because even where a particular nucleic acid molecule does not encode a polypeptide having BCSG1 activity, one of skill in the art would still know how to use the nucleic acid molecule, for instance, as a hybridization probe or a polymerase chain reaction (PCR) primer. Uses of the nucleic acid molecules of the present invention that do not encode a polypeptide having BCSG1 activity include, *inter alia*, (1) isolating the BCSG1 gene or allelic variants thereof in a cDNA library; (2) *in situ* hybridization (e.g., "FISH") to metaphase chromosomal spreads to provide precise chromosomal location of the BCSG1 gene, as described in Verma *et al.*, *Human Chromosomes: A Manual of Basic Techniques*, Pergamon Press, New York (1988); and Northern Blot analysis for detecting BCSG1 mRNA expression in specific tissues.

Preferred, however, are nucleic acid molecules having sequences at least 90%, 95%, 96%, 97%, 98% or 99% identical to the nucleic acid sequence shown in Figure 1 (SEQ ID NO:1) or to the nucleic acid sequence of the deposited

cDNA which do, in fact, encode a polypeptide having BCSG1 protein activity. By "a polypeptide having BCSG1 activity" is intended polypeptides exhibiting activity similar, but not necessarily identical, to an activity of the BCSG1 protein of the invention, as measured in a particular biological assay. BCSG1 protein is
5 believed to be involved with apoptosis. BCSG1 protein activity can be measured using assays that measure apoptosis. For example, human breast cancer cells cultured on Lab-Tek chamber slides (Nunc, Inc.) are treated with or without recombinant BCSG1 protein or a candidate BCSG1 protein. The cells are then treated with several concentrations of an apoptotic inducer, such as adriamycin.
10 Apoptosis is compared between the treated and control cells where DNA fragmentation is the criteria for apoptotic death using the following assay. At various time points after the adriamycin treatment, adherent cells are stained with DNA-specific fluorochrome diamino-2 phenylindole (Boehringer Mannheim) in a 1µg/ml methanol solution. Cells are counted within 20 minutes of staining on
15 a Zeiss Axiophot epifluoresence microscope. Experiments are performed in triplicate with at least 150 cells scored at each point. Fragmented or condensed nuclei are scored as apoptotic. Intact or mitotic nuclei are scored as normal.

Of course, due to the degeneracy of the genetic code, one of ordinary skill in the art will immediately recognize that a large number of the nucleic acid
20 molecules having a sequence at least 90%, 95%, 96%, 97%, 98%, or 99% identical to the nucleic acid sequence of the deposited cDNA or the nucleic acid sequence shown in Figure 1 (SEQ ID NO:1) will encode a polypeptide "having BCSG1 protein activity." In fact, since degenerate variants of these nucleotide sequences all encode the same polypeptide, this will be clear to the skilled artisan
25 even without performing the above described comparison assay. It will be further recognized in the art that, for such nucleic acid molecules that are not degenerate variants, a reasonable number will also encode a polypeptide having BCSG1 protein activity. This is because the skilled artisan is fully aware of amino acid substitutions that are either less likely or not likely to significantly effect protein

-14-

function (e.g., replacing one aliphatic amino acid with a second aliphatic amino acid).

For example, guidance concerning how to make phenotypically silent amino acid substitutions is provided in Bowie, J. U. *et al.*, "Deciphering the Message in Protein Sequences: Tolerance to Amino Acid Substitutions," *Science* 247:1306-1310 (1990), wherein the authors indicate that proteins are surprisingly tolerant of amino acid substitutions.

Vectors and Host Cells

The present invention also relates to vectors which include the isolated DNA molecules of the present invention, host cells which are genetically engineered with the recombinant vectors, and the production of BCSG1 polypeptides or fragments thereof by recombinant techniques.

The polynucleotides may be joined to a vector containing a selectable marker for propagation in a host. Generally, a plasmid vector is introduced in a precipitate, such as a calcium phosphate precipitate, or in a complex with a charged lipid. If the vector is a virus, it may be packaged *in vitro* using an appropriate packaging cell line and then transduced into host cells.

The DNA insert should be operatively linked to an appropriate promoter, such as the phage lambda PL promoter, the *E. coli lac*, *trp* and *tac* promoters, the SV40 early and late promoters and promoters of retroviral LTRs, to name a few. Other suitable promoters will be known to the skilled artisan. The expression constructs will further contain sites for transcription initiation, termination and, in the transcribed region, a ribosome binding site for translation. The coding portion of the mature transcripts expressed by the constructs will preferably include a translation initiating at the beginning and a termination codon (UAA, UGA or UAG) appropriately positioned at the end of the polypeptide to be translated.

-15-

As indicated, the expression vectors will preferably include at least one selectable marker. Such markers include dihydrofolate reductase or neomycin resistance for eukaryotic cell culture and tetracycline or ampicillin resistance genes for culturing in *E. coli* and other bacteria. Representative examples of appropriate hosts include, but are not limited to, bacterial cells, such as *E. coli*, *Streptomyces* and *Salmonella typhimurium* cells; fungal cells, such as yeast cells; insect cells such as *Drosophila* S2 and *Spodoptera* Sf9 cells; animal cells such as CHO, COS and Bowes melanoma cells; and plant cells. Appropriate culture mediums and conditions for the above-described host cells are known in the art.

In addition to the use of expression vectors in the practice of the present invention, the present invention further includes novel expression vectors comprising operator and promoter elements operatively linked to nucleotide sequences encoding a protein of interest. One example of such a vector is pHE4-5 which is described in detail below.

As summarized in Figures 3 and 4, components of the pHE4-5 vector (SEQ ID NO:10) include: 1) a neomycinphosphotransferase gene as a selection marker, 2) an *E. coli* origin of replication, 3) a T5 phage promoter sequence, 4) two *lac* operator sequences, 5) a Shine-Delgarno sequence, 6) the lactose operon repressor gene (*lacIq*). The origin of replication (*oriC*) is derived from pUC19 (LTI, Gaithersburg, MD). The promoter sequence and operator sequences were made synthetically. Synthetic production of nucleic acid sequences is well known in the art. CLONTECH 95/96 Catalog, pages 215-216, CLONTECH, 1020 East Meadow Circle, Palo Alto, CA 94303. A nucleotide sequence encoding BSCG-1 (SEQ ID NO:1), is operatively linked to the promoter and operator by inserting the nucleotide sequence between the NdeI and Asp718 sites of the pHE4-5 vector.

As noted above, the pHE4-5 vector contains a *lacIq* gene. *LacIq* is an allele of the *lacI* gene which confers tight regulation of the *lac* operator. Amann, E. et al., *Gene* 69:301-315 (1988); Stark, M., *Gene* 51:255-267 (1987). The *lacIq* gene encodes a repressor protein which binds to *lac* operator sequences and

blocks transcription of down-stream (*i.e.*, 3') sequences. However, the *lacIq* gene product dissociates from the *lac* operator in the presence of either lactose or certain lactose analogs, *e.g.*, isopropyl B-D-thiogalactopyranoside (IPTG). BSCG-1 thus is not produced in appreciable quantities in uninduced host cells
5 containing the pHE4-5 vector. Induction of these host cells by the addition of an agent such as IPTG, however, results in the expression of the BSCG-1 coding sequence.

The promoter/operator sequences of the pHE4-5 vector (SEQ ID NO:11) comprise a T5 phage promoter and two *lac* operator sequences. One operator is
10 located 5' to the transcriptional start site and the other is located 3' to the same site. These operators, when present in combination with the *lacIq* gene product, confer tight repression of down-stream sequences in the absence of a *lac* operon inducer, *e.g.*, IPTG. Expression of operatively linked sequences located down-stream from the *lac* operators may be induced by the addition of a *lac*
15 operon inducer, such as IPTG. Binding of a *lac* inducer to the *lacIq* proteins results in their release from the *lac* operator sequences and the initiation of transcription of operatively linked sequences. *Lac* operon regulation of gene expression is reviewed in Devlin, T., TEXTBOOK OF BIOCHEMISTRY WITH CLINICAL CORRELATIONS, 4th Edition (1997), pages 802-807.

The pHE4 series of vectors contain all of the components of the pHE4-5 vector except for the BSCG-1 coding sequence. Features of the pHE4 vectors include optimized synthetic T5 phage promoter, *lac* operator, and Shine-Delagarno sequences. Further, these sequences are also optimally spaced so that
20 expression of an inserted gene may be tightly regulated and high level of expression occurs upon induction.
25

Among known bacterial promoters suitable for use in the production of proteins of the present invention include the *E. coli lacI* and *lacZ* promoters, the T3 and T7 promoters, the *gpt* promoter, the lambda PR and PL promoters and the *trp* promoter. Suitable eukaryotic promoters include the CMV immediate early
30 promoter, the HSV thymidine kinase promoter, the early and late SV40

-17-

promoters, the promoters of retroviral LTRs, such as those of the Rous Sarcoma Virus (RSV), and metallothionein promoters, such as the mouse metallothionein-I promoter.

5 The pHE4-5 vector also contains a Shine-Delgarno sequence 5' to the AUG initiation codon. Shine-Delgarno sequences are short sequences generally located about 10 nucleotides up-stream (*i.e.*, 5') from the AUG initiation codon. These sequences essentially direct prokaryotic ribosomes to the AUG initiation codon.

10 Thus, the present invention is also directed to expression vector useful for the production of the proteins of the present invention. This aspect of the invention is exemplified by the pHE4-5 vector (SEQ ID NO:10).

15 Among vectors preferred for use in bacteria include pQE70, pQE60 and pQE-9, available from Qiagen; pBS vectors, Phagescript vectors, Bluescript vectors, pNH8A, pNH16a, pNH18A, pNH46A, available from Stratagene; and ptre99a, pKK223-3, pKK233-3, pDR540, pRIT5 available from Pharmacia. Among preferred eukaryotic vectors are pWLNEO, pSV2CAT, pOG44, pXT1 and pSG available from Stratagene; and pSVK3, pBPV, pMSG and pSVL available from Pharmacia. Other suitable vectors will be readily apparent to the skilled artisan.

20 Introduction of the construct into the host cell can be effected by calcium phosphate transfection, DEAE-dextran mediated transfection, cationic lipid-mediated transfection, electroporation, transduction, infection or other methods. Such methods are described in many standard laboratory manuals, such as Davis *et al.*, *Basic Methods In Molecular Biology* (1986).

25 The polypeptide may be expressed in a modified form, such as a fusion protein, and may include not only secretion signals, but also additional heterologous functional regions. For instance, a region of additional amino acids, particularly charged amino acids, may be added to the N-terminus of the polypeptide to improve stability and persistence in the host cell, during
30 purification, or during subsequent handling and storage. Also, peptide moieties

-18-

may be added to the polypeptide to facilitate purification. Such regions may be removed prior to final preparation of the polypeptide. The addition of peptide moieties to polypeptides to engender secretion or excretion, to improve stability and to facilitate purification, among others, are familiar and routine techniques in the art. A preferred fusion protein comprises a heterologous region from immunoglobulin that is useful to solubilize proteins. For example, EP-A-O 464 533 (Canadian counterpart 2045869) discloses fusion proteins comprising various portions of constant region of immunoglobulin molecules together with another human protein or part thereof. In many cases, the Fc part in a fusion protein is thoroughly advantageous for use in therapy and diagnosis and thus results, for example, in improved pharmacokinetic properties (EP-A 0232 262). On the other hand, for some uses it would be desirable to be able to delete the Fc part after the fusion protein has been expressed, detected and purified in the advantageous manner described. This is the case when Fc portion proves to be a hindrance to use in therapy and diagnosis, for example when the fusion protein is to be used as antigen for immunizations. In drug discovery, for example, human proteins, such as, hIL5-receptor has been fused with Fc portions for the purpose of high-throughput screening assays to identify antagonists of hIL-5. See, D. Bennett *et al.*, *Journal of Molecular Recognition*, Vol. 8:52-58 (1995) and K. Johanson *et al.*, *The Journal of Biological Chemistry*, Vol. 270, No. 16:9459-9471 (1995).

The BCSG1 protein can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography ("HPLC") is employed for purification. Polypeptides of the present invention include naturally purified products, products of chemical synthetic procedures, and products produced by recombinant techniques from a prokaryotic or

5 eukaryotic host, including, for example, bacterial, yeast, higher plant, insect and mammalian cells. Depending upon the host employed in a recombinant production procedure, the polypeptides of the present invention may be glycosylated or may be non-glycosylated. In addition, polypeptides of the invention may also include an initial modified methionine residue, in some cases as a result of host-mediated processes.

BCSG1 Polypeptides and Fragments

10 The invention further provides an isolated BCSG1 polypeptide having the amino acid sequence encoded by the deposited cDNA clones, or the amino acid sequence in Figure 1 (SEQ ID NO:2), or a peptide or polypeptide comprising a portion of the above polypeptides.

15 It will be recognized in the art that some amino acid sequences of the BCSG1 polypeptide can be varied without significant effect of the structure or function of the protein. If such differences in sequence are contemplated, it should be remembered that there will be critical areas on the protein which determine activity.

20 Thus, the invention further includes variations of the BCSG1 polypeptide which show substantial BCSG1 polypeptide activity or which include regions of BCSG1 protein such as the protein portions discussed below. Such mutants include deletions, insertions, inversions, repeats, and type substitutions. As indicated above, guidance concerning which amino acid changes are likely to be phenotypically silent can be found in Bowie, J.U., *et al.*, "Deciphering the Message in Protein Sequences: Tolerance to Amino Acid Substitutions," *Science* 247:1306-1310 (1990).

25 Thus, the fragment, derivative or analog of the polypeptide of Figure 1 (SEQ ID NO:2), or that encoded by the deposited cDNA, may be (i) one in which one or more of the amino acid residues are substituted with a conserved or non-conserved amino acid residue (preferably a conserved amino acid residue) and

-20-

such substituted amino acid residue may or may not be one encoded by the genetic code, or (ii) one in which one or more of the amino acid residues includes a substituent group, or (iii) one in which the polypeptide is fused with another compound, such as a compound to increase the half-life of the polypeptide (for example, polyethylene glycol), or (iv) one in which the additional amino acids are fused to the polypeptide, such as an IgG Fc fusion region peptide or leader or secretory sequence or a sequence which is employed for purification of the polypeptide or a proprotein sequence. Such fragments, derivatives and analogs are deemed to be within the scope of those skilled in the art from the teachings herein.

Of particular interest are substitutions of charged amino acids with another charged amino acid and with neutral or negatively charged amino acids. The latter results in proteins with reduced positive charge to improve the characteristics of the BCSG1 protein. The prevention of aggregation is highly desirable. Aggregation of proteins not only results in a loss of activity but can also be problematic when preparing pharmaceutical formulations, because they can be immunogenic. (Pinckard *et al.*, *Clin Exp. Immunol.* 2:331-340 (1967); Robbins *et al.*, *Diabetes* 36:838-845 (1987); Cleland *et al.* *Crit. Rev. Therapeutic Drug Carrier Systems* 10:307-377 (1993)).

As indicated, changes are preferably of a minor nature, such as conservative amino acid substitutions that do not significantly affect the folding or activity of the protein (see Table 1).

TABLE 1. Conservative Amino Acid Substitutions.

Aromatic	Phenylalanine Tryptophan Tyrosine
Hydrophobic	Leucine Isoleucine Valine
Polar	Glutamine Asparagine
Basic	Arginine Lysine Histidine
Acidic	Aspartic Acid Glutamic Acid
Small	Alanine Serine Threonine Methionine Glycine

Of course, the number of amino acid substitutions a skilled artisan would make depends on many factors, including those described above. Generally speaking, the number of substitutions for any given AIM-II polypeptide will not be more than 50, 40, 30, 25, 20, 15, 10, 5 or 3.

Amino acids in the BCSG1 protein of the present invention that are essential for function can be identified by methods known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham and Wells, *Science* 244:1081-1085 (1989)). The latter procedure introduces single alanine mutations at every residue in the molecule. The resulting mutant molecules are then tested for biological activity such as receptor binding or *in vitro*, or *in vitro* proliferative activity. Sites that are critical for ligand-receptor binding can also be determined by structural analysis such as crystallization, nuclear magnetic resonance or photoaffinity labeling (Smith *et al.*, *J. Mol. Biol.* 224:899-904 (1992) and de Vos *et al.* *Science* 255:306-312 (1992)).

-22-

The polypeptides of the present invention are preferably provided in an isolated form. By "isolated polypeptide" is intended a polypeptide removed from its native environment. Thus, a polypeptide produced and/or contained within a recombinant host cell is considered isolated for purposes of the present invention. Also intended as an "isolated polypeptide" are polypeptides that have been purified, partially or substantially, from a recombinant host cell or from a native source. For example, a recombinantly produced version of the BCSG1 polypeptide can be substantially purified by the one-step method described in Smith and Johnson, *Gene* 67:31-40 (1988).

The polypeptides of the present invention include the polypeptide encoded by the deposited cDNA; a polypeptide comprising amino acids about 1 to about 127 in SEQ ID NO:2 (Figure 1); a polypeptide comprising amino acids about 2 to about 127 in SEQ ID NO:2; as well as polypeptides which are at least 80% identical, more preferably at least 90% or 95% identical, still more preferably at least 96%, 97%, 98% or 99% identical to the polypeptide encoded by the deposited cDNA, to the polypeptide of Figure 1 (SEQ ID NO:2), and also include portions of such polypeptides with at least 30 amino acids and more preferably at least 50 amino acids.

By a polypeptide having an amino acid sequence at least, for example, 95% "identical" to a reference amino acid sequence of a BCSG1 polypeptide is intended that the amino acid sequence of the polypeptide is identical to the reference sequence except that the polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the reference amino acid of the BCSG1 polypeptide. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a reference amino acid sequence, up to 5% of the amino acid residues in the reference sequence may be deleted or substituted with another amino acid, or a number of amino acids up to 5% of the total amino acid residues in the reference sequence may be inserted into the reference sequence. These alterations of the reference sequence may occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere

-23-

between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

5 As a practical matter, whether any particular polypeptide is at least 90%, 95%, 96%, 97%, 98% or 99% identical to, for instance, the amino acid sequence shown in Figure 1 (SEQ ID NO:2) or to the amino acid sequence encoded by deposited cDNA clone can be determined conventionally using known computer programs such the Bestfit program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, 575
10 Science Drive, Madison, WI 53711. When using Bestfit or any other sequence alignment program to determine whether a particular sequence is, for instance, 95% identical to a reference sequence according to the present invention, the parameters are set, of course, such that the percentage of identity is calculated over the full length of the reference amino acid sequence and that gaps in
15 homology of up to 5% of the total number of amino acid residues in the reference sequence are allowed.

The polypeptide of the present invention could be used as a molecular weight marker on SDS-PAGE gels or on molecular sieve gel filtration columns using methods well known to those of skill in the art.

20 In another aspect, the invention provides a peptide or polypeptide comprising an epitope-bearing portion of a polypeptide of the invention. The epitope of this polypeptide portion is an immunogenic or antigenic epitope of a polypeptide described herein. An "immunogenic epitope" is defined as a part of a protein that elicits an antibody response when the whole protein is the
25 immunogen. On the other hand, a region of a protein molecule to which an antibody can bind is defined as an "antigenic epitope." The number of immunogenic epitopes of a protein generally is less than the number of antigenic epitopes. See, for instance, Geysen *et al.*, *Proc. Natl. Acad. Sci. USA* 81:3998-4002 (1983).

As to the selection of peptides or polypeptides bearing an antigenic epitope (i.e., that contain a region of a protein molecule to which an antibody can bind), it is well known in that art that relatively short synthetic peptides that mimic part of a protein sequence are routinely capable of eliciting an antiserum that reacts with the partially mimicked protein. See, for instance, Sutcliffe, J. G., Shinnick, T. M., Green, N. and Learner, R.A. (1983) Antibodies that react with predetermined sites on proteins. *Science* 219:660-666. Peptides capable of eliciting protein-reactive sera are frequently represented in the primary sequence of a protein, can be characterized by a set of simple chemical rules, and are confined neither to immunodominant regions of intact proteins (i.e., immunogenic epitopes) nor to the amino or carboxyl terminals.

Antigenic epitope-bearing peptides and polypeptides of the invention are therefore useful to raise antibodies, including monoclonal antibodies, that bind specifically to a polypeptide of the invention. See, for instance, Wilson *et al.*, *Cell* 37:767-778 (1984) at 777.

Antigenic epitope-bearing peptides and polypeptides of the invention preferably contain a sequence of at least seven, more preferably at least nine and most preferably between about at least about 15 to about 30 amino acids contained within the amino acid sequence of a polypeptide of the invention.

Non-limiting examples of antigenic polypeptides or peptides that can be used to generate BCSG1-specific antibodies include: a polypeptide comprising amino acid residues from about 94 to about 107 in Figure 1 (SEQ ID NO:2); a polypeptide comprising amino acid residues from about 120 to about 127 in Figure 1 (SEQ ID NO:2). As indicated above, the inventors have determined that the above polypeptide fragments are antigenic regions of the BCSG1 protein.

The epitope-bearing peptides and polypeptides of the invention may be produced by any conventional means. Houghten, R. A. (1985) General method for the rapid solid-phase synthesis of large numbers of peptides: specificity of antigen-antibody interaction at the level of individual amino acids. *Proc. Natl. Acad. Sci. USA* 82:5131-5135. This "Simultaneous Multiple Peptide Synthesis

(SMPS)" process is further described in U.S. Patent No. 4,631,211 to Houghten *et al.* (1986).

As one of skill in the art will appreciate, BCSG1 polypeptides of the present invention and the epitope-bearing fragments thereof described above can be combined with parts of the constant domain of immunoglobulins (IgG), resulting in chimeric polypeptides. These fusion proteins facilitate purification and show an increased half-life *in vivo*. This has been shown, e.g., for chimeric proteins consisting of the first two domains of the human CD4-polypeptide and various domains of the constant regions of the heavy or light chains of mammalian immunoglobulins (EPA 394,827; Traunecker *et al.*, *Nature* 331:84-86 (1988)). Fusion proteins that have a disulfide-linked dimeric structure due to the IgG part can also be more efficient in binding and neutralizing other molecules than the monomeric BCSG1 protein or protein fragment alone (Fountoulakis *et al.*, *J. Biochem* 270:3958-3964 (1995)).

Cancer Diagnosis and Prognosis

There are two classes of genes affecting tumor development. Genes influencing the cancer phenotype that act directly as a result of changes (e.g., mutation) at the DNA level, such as BRCA1, BRCA2, and p53, are called Class I genes. The Class II genes affect the phenotype by modulation at the expression level. Development of breast cancer and subsequent malignant progression is associated with alterations of a variety of genes of both classes. Identification of quantitative changes in gene expression that occur in the malignant mammary gland, if sufficiently characterized, may yield novel molecular markers which may be useful in the diagnosis and treatment of human breast cancer.

The present inventors have identified a new breast cancer marker that is overexpressed in advanced infiltrating breast cancer cells. The lack of expression of BCSG1 in normal or benign breast epithelial cells and a weak expression in low grade *in situ* carcinomas suggest that overexpression of BCSG1 indicates

breast cancer malignant progression. (See, Examples 6 and 7). It is unlikely that BCSG1 is overexpressed as a secondary effect of cellular proliferation because no detectable BCSG1 expression is evident in rapidly proliferating nonmalignant breast lesions. (See, Example 7).

5 BCSG1 may be useful in clinical management and treatment of breast cancer. In this regard, the expression of BCSG1 transcripts was observed in the neoplastic epithelial cells of infiltrating breast carcinoma but not in epithelial cells of normal and benign breast tissue. (See, Example 7). The overexpression of BCSG1 in malignant infiltrating breast epithelial cells compared to the low
10 level expression in the low grade *in situ* carcinoma suggests that up-regulation of BCSG1 expression is associated with breast malignant progression and may signal the more advanced invasive/metastatic phenotype of human breast cancer. This implication is further supported by detection of BCSG1 expression in 4/4 breast cancer cell lines derived from ductal infiltrating carcinomas but not (0/3)
15 in breast cancer cell lines derived from primary solid carcinoma (See, Example 6). BCSG1 overexpression in ductal carcinoma *in situ* (DCIS) may indicate a malignant progression leading to metastasis. There was a marked increase in DCIS incidence beginning in the early 1980s (Emster, V.L., *et al.*, *JAMA* 275:913-918 (1996)). The total estimated number of DCIS cases in the United
20 States in 1992 was 200% higher than expected based on 1983 rates and trends between 1973 and 1983 (Emster, V.L., *et al.*, *JAMA* 275:913-918 (1996)). While early detection of invasive breast cancer is beneficial, the value of DCIS detection is currently unknown. There is cause for concern about the large number of DCIS cases that are being diagnosed as a consequence of screening mammography,
25 most of which are treated by some form of surgery. In addition, the proportion of cases treated by mastectomy may be inappropriately high (Emster, V.L., *et al.*, *JAMA* 275:913-918 (1996)). BCSG1 expression may provide some prognostic information on distinguishing the DCIS which is not likely to become invasive from the DCIS which is most likely to become invasive, which will help to reduce
30 some inappropriate or unnecessary mastectomies. In addition, the use of BCSG1

gene could be of great importance in differentiating atypical proliferative breast lesions from cancer and may be useful in screening of breast biopsies for potential abnormalities.

It is interesting to note that the predicted amino acid sequence of BCSG1 gene shares high sequence homology with the recently cloned non-A β component of Alzheimer's disease (AD) amyloid precursor protein (Ueda, K., *et al.*, *Proc. Natl. Acad. Sci. USA.* 90(23):11282-6 (1993)). A neuropathological hallmark of AD is a widespread amyloid deposition resulting from beta-amyloid precursor proteins (beta APPS). Beta APPs are large membrane-spanning proteins that either give rise to the beta A4 peptide (AB fragment) (Masters, C.L., *et al.*, *Proc. Natl. Acad. Sci. USA* 82:4245-4249 (1985)) or a non-A β component of AD amyloid (Ueda, K., *et al.*, *Proc. Natl. Acad. Sci. USA.* 90(23):11282-6 (1993)) that is either deposited in AD amyloid plaques or yielding soluble forms. While the insoluble membrane-bound AD amyloid destabilizes calcium homeostasis and thus renders cell vulnerable to excitotoxic conditions of calcium influx resulting from energy deprivation or overexcitation (Mattson, M.P., *et al.*, *Ann. N.Y. Acad. Sci.* 679:121 (1993)), the soluble AD amyloid proteins are neuroprotective against glucose deprivation and glutamate toxicity, perhaps through their ability to lower the intraneuronal calcium concentration (Barger, S.W., *J. Neurochem.* 64:2087-96 (1995)). It is possible that BCSG1, like soluble AD amyloid, may be potentially involved in tissue damage resulting from tissue remodeling due to the local cancer invasion. Nevertheless, Examples 6 and 7 demonstrate a stage-specific BCSG1 expression and an association of BCSG1 overexpression with clinical aggressiveness of breast cancers. BCSG1 overexpression may indicate breast cancer malignant progression from benign breast or low grade *in situ* carcinoma to the highly infiltrating carcinoma.

The Examples demonstrate that certain tissues in mammals with cancer express significantly enhanced levels of the BCSG1 protein and mRNA encoding the BCSG1 protein when compared to a corresponding "standard" mammal, i.e., a mammal of the same species not having the cancer. Further, it is believed that

-28-

enhanced levels of the BCSG1 protein can be detected in certain body fluids (e.g., sera, plasma, urine, and spinal fluid) from mammals with cancer when compared to sera from mammals of the same species not having the cancer. Thus, the invention provides a diagnostic method useful during tumor diagnosis, which involves assaying the expression level of the gene encoding the BCSG1 protein in mammalian cells or body fluid and comparing the gene expression level with a standard BCSG1 gene expression level, whereby an increase in the gene expression level over the standard is indicative of certain tumors.

Where a tumor diagnosis has already been made according to conventional methods, the present invention is useful as a prognostic indicator, whereby patients exhibiting enhanced BCSG1 gene expression will experience a worse clinical outcome relative to patients expressing the gene at a lower level.

By "assaying the expression level of the gene encoding the BCSG1 protein" is intended qualitatively or quantitatively measuring or estimating the level of the BCSG1 protein or the level of the mRNA encoding the BCSG1 protein in a first biological sample either directly (e.g., by determining or estimating absolute protein level or mRNA level) or relatively (e.g., by comparing to the BCSG1 protein level or mRNA level in a second biological sample).

Preferably, the BCSG1 protein level or mRNA level in the first biological sample is measured or estimated and compared to a standard BCSG1 protein level or mRNA level, the standard being taken from a second biological sample obtained from an individual not having the cancer. As will be appreciated in the art, once a standard BCSG1 protein level or mRNA level is known, it can be used repeatedly as a standard for comparison.

By "biological sample" is intended any biological sample obtained from an individual, cell line, tissue culture, or other source which contains BCSG1 protein or mRNA. Biological samples include mammalian body fluids (such as sera, plasma, urine, synovial fluid and spinal fluid) which contain secreted mature BCSG1 protein, and ovarian, prostate, heart, placenta, pancreas liver, spleen, lung, breast and umbilical tissue.

The present invention is useful for detecting cancer in mammals. In particular the invention is useful during diagnosis of the of following types of cancers in mammals: breast, ovarian, prostate, bone, liver, lung, pancreatic, and splenic. Preferred mammals include monkeys, apes, cats, dogs, cows, pigs, horses, rabbits and humans. Particularly preferred are humans.

Total cellular RNA can be isolated from a biological sample using the single-step guanidinium-thiocyanate-phenol-chloroform method described in Chomczynski and Sacchi, *Anal. Biochem.* 162:156-159 (1987). Levels of mRNA encoding the BCSG1 protein are then assayed using any appropriate method. These include Northern blot analysis (Harada *et al.*, *Cell* 63:303-312 (1990)), S1 nuclease mapping (Fujita *et al.*, *Cell* 49:357- 367 (1987)), the polymerase chain reaction (PCR), reverse transcription in combination with the polymerase chain reaction (RT-PCR) (Makino *et al.*, *Technique* 2:295-301 (1990)), and reverse transcription in combination with the ligase chain reaction (RT-LCR).

Assaying BCSG1 protein levels in a biological sample can occur using antibody-based techniques. For example, BCSG1 protein expression in tissues can be studied with classical immunohistological methods (Jalkanen, M., *et al.*, *J. Cell. Biol.* 101:976-985 (1985); Jalkanen, M., *et al.*, *J. Cell . Biol.* 105:3087-3096 (1987)).

Other antibody-based methods useful for detecting BCSG1 protein gene expression include immunoassays, such as the enzyme linked immunosorbent assay (ELISA) and the radioimmunoassay (RIA).

Suitable labels are known in the art and include enzyme labels, such as, Glucose oxidase, and radioisotopes, such as iodine (^{125}I , ^{121}I), carbon (^{14}C), sulfur (^{35}S), tritium (^3H), indium (^{112}In), and technetium ($^{99\text{m}}\text{Tc}$), and fluorescent labels, such as fluorescein and rhodamine, and biotin.

Chromosome Assays

The nucleic acid molecules of the present invention are also valuable for chromosome identification. The sequence is specifically targeted to and can hybridize with a particular location on an individual human chromosome. The mapping of DNAs to chromosomes according to the present invention is an important first step in correlating those sequences with genes associated with disease.

In certain preferred embodiments in this regard, the cDNA herein disclosed is used to clone genomic DNA of a BCSG1 protein gene. This can be accomplished using a variety of well known techniques and libraries, which generally are available commercially. The genomic DNA then is used for *in situ* chromosome mapping using well known techniques for this purpose.

In addition, in some cases, sequences can be mapped to chromosomes by preparing PCR primers (preferably 15-25 bp) from the cDNA. Computer analysis of the 3' untranslated region of the gene is used to rapidly select primers that do not span more than one exon in the genomic DNA, thus complicating the amplification process. These primers are then used for PCR screening of somatic cell hybrids containing individual human chromosomes.

Fluorescence *in situ* hybridization ("FISH") of a cDNA clone to a metaphase chromosomal spread can be used to provide a precise chromosomal location in one step. This technique can be used with probes from the cDNA as short as 50 or 60 bp. For a review of this technique, see Verma *et al.*, *Human Chromosomes: A Manual Of Basic Techniques*, Pergamon Press, New York (1988).

Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. Such data are found, for example, in V. McKusick, *Mendelian Inheritance In Man*, available on-line through Johns Hopkins University, Welch Medical Library. The relationship between genes and diseases that have been

-31-

mapped to the same chromosomal region are then identified through linkage analysis (coinheritance of physically adjacent genes).

Next, it is necessary to determine the differences in the cDNA or genomic sequence between affected and unaffected individuals. If a mutation is observed in some or all of the affected individuals but not in any normal individuals, then the mutation is likely to be the causative agent of the disease.

Having generally described the invention, the same will be more readily understood by reference to the following examples, which are provided by way of illustration and are not intended as limiting.

Examples

Example 1: Expression and Purification of BCSG1 in E. coli

The bacterial expression vector pQE9 (pD10) is used for bacterial expression in this example. (QIAGEN, Inc., 9259 Eton Avenue, Chatsworth, CA, 91311). pQE9 encodes ampicillin antibiotic resistance ("Amp") and contains a bacterial origin of replication ("ori"), an IPTG inducible promoter, a ribosome binding site ("RBS"), six codons encoding histidine residues that allow affinity purification using nickel-nitrilo-tri-acetic acid ("Ni-NTA") affinity resin sold by QIAGEN, Inc., *supra*, and suitable single restriction enzyme cleavage sites. These elements are arranged such that an inserted DNA fragment encoding a polypeptide expresses that polypeptide with the six His residues (i.e., a "6 X His tag")) covalently linked to the amino terminus of that polypeptide.

The DNA sequence encoding the desired portion BCSG1 protein sequence is amplified from the deposited cDNA clone using PCR oligonucleotide primers which anneal to the amino terminal sequences of the desired portion of the BCSG1 protein and to sequences in the deposited construct 3' to the cDNA coding sequence. Additional nucleotides containing restriction sites to facilitate

-32-

cloning in the pQE9 vector are added to the 5' and 3' primer sequences, respectively.

For cloning the mature protein, the 5' primer has the sequence 5' GGGGATCCATGTTTCAAGAAGG 3' (SEQ ID NO:3) containing the underlined BamHI restriction site followed by 16 nucleotides complementary to the amino terminal coding sequence of the BCSG1 sequence in Figure 1. One of ordinary skill in the art would appreciate, of course, that the point in the protein coding sequence where the 5' primer begins may be varied to amplify a DNA segment encoding any desired portion of the complete BCSG1 protein shorter or longer than the protein. The 3' primer has the sequence 5' GGAAGCTTCTAGTCTCCCCCACTCTGG 3' (SEQ ID NO:4) containing the underlined HindIII restriction site followed by 19 nucleotides complementary to the non-coding sequence of the BCSG1 DNA sequence in Figure 1.

The amplified BCSG1 DNA fragment and the vector pQE9 are digested with BamHI/HindIII and the digested DNAs are then ligated together. Insertion of the BCSG1 DNA into the restricted pQE9 vector places the BCSG1 protein coding region downstream from the IPTG-inducible promoter and in-frame with an initiating AUG and the six histidine codons.

The ligation mixture is transformed into competent *E. coli* cells using standard procedures such as those described in Sambrook *et al.*, *Molecular Cloning: a Laboratory Manual, 2nd Ed.*; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY (1989). *E. coli* strain M15/rep4, containing multiple copies of the plasmid pREP4, which expresses the lac repressor and confers kanamycin resistance ("Kan^r"), is used in carrying out the illustrative example described herein. This strain, which is only one of many that are suitable for expressing BCSG1 protein, is available commercially from QIAGEN, Inc., *supra*. Transformants are identified by their ability to grow on LB plates in the presence of ampicillin and kanamycin. Plasmid DNA is isolated from resistant colonies and the identity of the cloned DNA confirmed by restriction analysis, PCR and DNA sequencing.

Clones containing the desired constructs are grown overnight ("O/N") in liquid culture in LB media supplemented with both ampicillin (100 µg/ml) and kanamycin (25 µg/ml). The O/N culture is used to inoculate a large culture, at a dilution of approximately 1:25 to 1:250. The cells are grown to an optical density at 600 nm ("OD600") of between 0.4 and 0.6. Isopropyl- β -D-thiogalactopyranoside ("IPTG") is then added to a final concentration of 1 mM to induce transcription from the lac repressor sensitive promoter, by inactivating the lacI repressor. Cells subsequently are incubated further for 3 to 4 hours. Cells then are harvested by centrifugation.

The cells are then stirred for 3-4 hours at 4°C in 6 M guanidine-HCl, pH 8. The cell debris is removed by centrifugation, and the supernatant containing the BCSG1 is loaded onto a nickel-nitrilo-tri-acetic acid ("NiNTA") affinity resin column (available from QIAGEN, Inc., *supra*). Proteins with a 6 x His tag bind to the Ni-NTA resin with high affinity and can be purified in a simple one-step procedure (for details see: The QIAexpressionist, 1995, QIAGEN, Inc., *supra*). Briefly the supernatant is loaded onto the column in 6 M guanidine-HCl, pH 8, the column is first washed with 10 volumes of 6 M guanidine-HCl, pH 8, then washed with 10 volumes of 6 M guanidine-HCl pH 6, and finally the BCSG1 is eluted with 6 M guanidine-HCl, pH 5.

The purified protein is then renatured by dialyzing it against phosphate-buffered saline (PBS) or 50 mM Na-acetate, pH 6 buffer plus 200 mM NaCl. Alternatively, the protein can be successfully refolded while immobilized on the Ni-NTA column. The recommended conditions are as follows: renature using a linear 6M-1M urea gradient in 500 mM NaCl, 20% glycerol, 20 mM Tris/HCl pH 7.4, containing protease inhibitors. The renaturation should be performed over a period of 1.5 hours or more. After renaturation the proteins can be eluted by the addition of 250 mM imidazole. Imidazole is removed by a final dialyzing step against PBS or 50 mM sodium acetate pH 6 buffer plus 200 mM NaCl. The purified protein is stored at 4°C or frozen at -80°C.

Example 2: Cloning and Expression of BCSG1 protein in a Baculovirus Expression System

In this illustrative example, the plasmid shuttle vector pA2 GP is used to insert the cloned DNA encoding the protein into a baculovirus to express the BCSG1 protein, using a baculovirus leader and standard methods as described in Summers *et al.*, *A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures*, Texas Agricultural Experimental Station Bulletin No. 1555 (1987). This expression vector contains the strong polyhedrin promoter of the *Autographa californica* nuclear polyhedrosis virus (AcMNPV) followed by the secretory signal peptide (leader) of the baculovirus gp67 protein and convenient restriction sites such as BamHI, Xba I and Asp718. The polyadenylation site of the simian virus 40 ("SV40") is used for efficient polyadenylation. For easy selection of recombinant virus, the plasmid contains the beta-galactosidase gene from *E. coli* under control of a weak *Drosophila* promoter in the same orientation, followed by the polyadenylation signal of the polyhedrin gene. The inserted genes are flanked on both sides by viral sequences for cell-mediated homologous recombination with wild-type viral DNA to generate viable virus that expresses the cloned polynucleotide.

Many other baculovirus vectors could be used in place of the vector above, such as pAc373, pVL941 and pAcIM1, as one skilled in the art would readily appreciate, as long as the construct provides appropriately located signals for transcription, translation, secretion and the like, including a signal peptide and an in-frame AUG as required. Such vectors are described, for instance, in Luckow *et al.*, *Virology* 170:31-39.

The cDNA sequence encoding the BCSG1 protein in the deposited clone shown in Figure 1 (SEQ ID NO:2), is amplified using PCR oligonucleotide primers corresponding to the 5' and 3' sequences of the gene.

The 5' primer has the sequence 5' GGGGATCCcGATGTTTTCAAGAAGG 3' (SEQ ID NO:5) (the lowercase "c"

is a nucleotide included to preserve the coding frame) containing the underlined BamHI restriction enzyme site followed by 16 bases of the sequence of the BCSG1 protein shown in Figure 1, beginning with the N-terminus of the protein. The 3' primer has the sequence 5'GGGGTACCCTAGTCTCCCCCACTCTGG 3' (SEQ ID NO:6) containing the underlined Asp718 restriction site followed by 18 nucleotides complementary to the 3' noncoding sequence in Figure 1.

The amplified fragment is isolated from a 1% agarose gel using a commercially available kit ("GeneClean," BIO 101 Inc., La Jolla, Ca.). The fragment then is digested with BamHI/Asp718 and again is purified on a 1% agarose gel. This fragment is designated herein "F1".

The plasmid is digested with the restriction enzymes BamHI/Asp718 and optionally, can be dephosphorylated using calf intestinal phosphatase, using routine procedures known in the art. The DNA is then isolated from a 1% agarose gel using a commercially available kit ("GeneClean" BIO 101 Inc., La Jolla, Ca.). This vector DNA is designated herein "V1".

Fragment F1 and the dephosphorylated plasmid V1 are ligated together with T4 DNA ligase. *E. coli* HB101 or other suitable *E. coli* hosts such as XL-1 Blue (Stratagene Cloning Systems, La Jolla, CA) cells are transformed with the ligation mixture and spread on culture plates. Bacteria are identified that contain the plasmid with the human BCSG1 gene using the PCR method, in which one of the primers that is used to amplify the gene and the second primer is from well within the vector so that only those bacterial colonies containing the BCSG1 gene fragment will show amplification of the DNA. The sequence of the cloned fragment is confirmed by DNA sequencing. This plasmid is designated herein pBac BCSG1.

Five µg of the plasmid pBacBCSG1 is co-transfected with 1.0 µg of a commercially available linearized baculovirus DNA ("BaculoGold™ baculovirus DNA", Pharmingen, San Diego, CA.), using the lipofection method described by Felgner *et al.*, *Proc. Natl. Acad. Sci. USA* 84:7413-7417 (1987). 1 µg of BaculoGold™ virus DNA and 5 µg of the plasmid pBac BCSG1 are mixed in a

sterile well of a microtiter plate containing 50 μ l of serum-free Grace's medium (Life Technologies Inc., Gaithersburg, MD). Afterwards, 10 μ l Lipofectin plus 90 μ l Grace's medium are added, mixed and incubated for 15 minutes at room temperature. Then the transfection mixture is added drop-wise to Sf9 insect cells (ATCC CRL 1711) seeded in a 35 mm tissue culture plate with 1 ml Grace's medium without serum. The plate is rocked back and forth to mix the newly added solution. The plate is then incubated for 5 hours at 27°C. After 5 hours the transfection solution is removed from the plate and 1 ml of Grace's insect medium supplemented with 10% fetal calf serum is added. The plate is put back into an incubator and cultivation is continued at 27°C for four days.

After four days the supernatant is collected and a plaque assay is performed, as described by Summers and Smith, *supra*. An agarose gel with "Blue Gal" (Life Technologies Inc., Gaithersburg) is used to allow easy identification and isolation of gal-expressing clones, which produce blue-stained plaques. (A detailed description of a "plaque assay" of this type can also be found in the user's guide for insect cell culture and baculovirology distributed by Life Technologies Inc., Gaithersburg, page 9-10). After appropriate incubation, blue stained plaques are picked with the tip of a micropipettor (e.g., Eppendorf). The agar containing the recombinant viruses is then resuspended in a microcentrifuge tube containing 200 μ l of Grace's medium and the suspension containing the recombinant baculovirus is used to infect Sf9 cells seeded in 35 mm dishes. Four days later the supernatants of these culture dishes are harvested and then they are stored at 4°C. The recombinant virus is called V-BCSG1.

To verify the expression of the BCSG1 gene, Sf9 cells are grown in Grace's medium supplemented with 10% heat inactivated FBS. The cells are infected with the recombinant baculovirus V-BCSG1 at a multiplicity of infection ("MOI") of about 2. Six hours later the medium is removed and is replaced with SF900 II medium minus methionine and cysteine (available from Life Technologies Inc., Rockville, MD). If radiolabeled proteins are desired, 42 hours later, 5 μ Ci of 35 S-methionine and 5 μ Ci 35 S-cysteine (available from Amersham)

-37-

are added. The cells are further incubated for 16 hours and then they are harvested by centrifugation. The proteins in the supernatant as well as the intracellular proteins are analyzed by SDS-PAGE followed by autoradiography (if radiolabeled). Microsequencing of the amino acid sequence of the amino terminus of purified protein may be used to determine the amino terminal sequence of the mature protein and thus the cleavage point and length of the secretory signal peptide.

Example 3: Cloning and Expression of BCSG1 in Mammalian Cells

A typical mammalian expression vector contains the promoter element, which mediates the initiation of transcription of mRNA, the protein coding sequence, and signals required for the termination of transcription and polyadenylation of the transcript. Additional elements include enhancers, Kozak sequences and intervening sequences flanked by donor and acceptor sites for RNA splicing. Highly efficient transcription can be achieved with the early and late promoters from SV40, the long terminal repeats (LTRs) from Retroviruses, e.g., RSV, HTLV I, HIV I and the early promoter of the cytomegalovirus (CMV). However, cellular elements can also be used (e.g., the human actin promoter). Suitable expression vectors for use in practicing the present invention include, for example, vectors such as PSVL and PMSG (Pharmacia, Uppsala, Sweden), pRSVcat (ATCC 37152), pSV2dhfr (ATCC 37146) and pBC12MI (ATCC 67109). Mammalian host cells that could be used include, human Hela 293, H9 and Jurkat cells, mouse NIH3T3 and C127 cells, Cos 1, Cos 7 and CV 1, quail QC1-3 cells, mouse L cells and Chinese hamster ovary (CHO) cells.

Alternatively, the gene can be expressed in stable cell lines that contain the gene integrated into a chromosome. The co-transfection with a selectable marker such as dhfr, gpt, neomycin, or hygromycin allows the identification and isolation of the transfected cells.

5 The transfected gene can also be amplified to express large amounts of the encoded protein. The DHFR (dihydrofolate reductase) marker is useful to develop cell lines that carry several hundred or even several thousand copies of the gene of interest. Another useful selection marker is the enzyme glutamine synthase (GS) (Murphy *et al.*, *Biochem J.* 227:277-279 (1991); Bebbington *et al.*, *Bio/Technology* 10:169-175 (1992)). Using these markers, the mammalian cells are grown in selective medium and the cells with the highest resistance are selected. These cell lines contain the amplified gene(s) integrated into a chromosome. Chinese hamster ovary (CHO) and NSO cells are often used for the production of proteins.

10 The expression vectors pC1 and pC4 contain the strong promoter (LTR) of the Rous Sarcoma Virus (Cullen *et al.*, *Molecular and Cellular Biology*, 438-447 (March, 1985)) plus a fragment of the CMV-enhancer (Boshart *et al.*, *Cell* 41:521-530 (1985)). Multiple cloning sites, e.g., with the restriction enzyme cleavage sites BamHI, XbaI and Asp718, facilitate the cloning of the gene of interest. The vectors contain in addition the 3' intron, the polyadenylation and termination signal of the rat preproinsulin gene.

Example 3(a): Cloning and Expression in COS Cells

20 The expression plasmid, pBCSG1 HA, is made by cloning a cDNA encoding BCSG1 into the expression vector pcDNAI/Amp or pcDNAIII (which can be obtained from Invitrogen, Inc.).

25 The expression vector pcDNAI/amp contains: (1) an *E. coli* origin of replication effective for propagation in *E. coli* and other prokaryotic cells; (2) an ampicillin resistance gene for selection of plasmid-containing prokaryotic cells; (3) an SV40 origin of replication for propagation in eukaryotic cells; (4) a CMV promoter, a polylinker, an SV40 intron; (5) several codons encoding a hemagglutinin fragment (i.e., an "HA" tag to facilitate purification) followed by a termination codon and polyadenylation signal arranged so that a cDNA can be

conveniently placed under expression control of the CMV promoter and operably linked to the SV40 intron and the polyadenylation signal by means of restriction sites in the polylinker. The HA tag corresponds to an epitope derived from the influenza hemagglutinin protein described by Wilson *et al.*, *Cell* 37:767 (1984).
5 The fusion of the HA tag to the target protein allows easy detection and recovery of the recombinant protein with an antibody that recognizes the HA epitope. pcDNAIII contains, in addition, the selectable neomycin marker.

A DNA fragment encoding the BCSG1 is cloned into the polylinker region of the vector so that recombinant protein expression is directed by the
10 CMV promoter. The plasmid construction strategy is as follows. The BCSG1 cDNA of the deposited clone is amplified using primers that contain convenient restriction sites, much as described above for construction of vectors for expression of BCSG1 in *E. coli*. Suitable primers include the following, which are used in this example. The 5' primer, containing the underlined BamHI site,
15 a Kozak sequence, an AUG start codon and 4 codons of the 5' coding region of the complete BCSG1 has the following sequence: 5' GGGGATccgccaccATGTTTCAAGAAGG 3' (SEQ ID NO:7) (Kozak sequence is represented by the lowercase letters). The 3' primer, containing the underlined BamHI site, a stop codon, and 19 bp of 3' coding sequence has the following
20 sequence (at the 3' end): 5' GGGGATCCTCagaagcgtagtctgggacgtcgatgggtaCTAGTCTCCCCCACTCTGG 3' (SEQ ID NO:8) (the HA tag is represented by the lowercase letters).

The PCR amplified DNA fragment and the vector, pcDNAI/Amp, are digested with BamHI and then ligated. The ligation mixture is transformed into
25 *E. coli* strain SURE (available from Stratagene Cloning Systems, 11099 North Torrey Pines Road, La Jolla, CA 92037), and the transformed culture is plated on ampicillin media plates which then are incubated to allow growth of ampicillin resistant colonies. Plasmid DNA is isolated from resistant colonies and examined by restriction analysis or other means for the presence of the BCSG1-encoding
30 fragment.

For expression of recombinant BCSG1, COS cells are transfected with an expression vector, as described above, using DEAE-DEXTRAN, as described, for instance, in Sambrook *et al.*, *Molecular Cloning: a Laboratory Manual*, Cold Spring Laboratory Press, Cold Spring Harbor, New York (1989). Cells are

5 incubated under conditions for expression of BCSG1 by the vector.

Expression of the BCSG1-HA fusion protein is detected by radiolabeling and immunoprecipitation, using methods described in, for example Harlow *et al.*, *Antibodies: A Laboratory Manual, 2nd Ed.*; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1988). To this end, two days after

10 transfection, the cells are labeled by incubation in media containing ³⁵S-cysteine for 8 hours. The cells and the media are collected, and the cells are washed and lysed with detergent-containing RIPA buffer: 150 mM NaCl, 1% NP-40, 0.1% SDS, 0.5% DOC, 50 mM TRIS, pH 7.5, as described by Wilson *et al.* cited above. Proteins are precipitated from the cell lysate and from the culture media

15 using an HA-specific monoclonal antibody. The precipitated proteins then are analyzed by SDS-PAGE and autoradiography. An expression product of the expected size is seen in the cell lysate, which is not seen in negative controls.

Example 3(b): Cloning and Expression in CHO Cells

The vector pC4 is used for the expression of BCSG1 protein. Plasmid

20 pC4 is a derivative of the plasmid pSV2-dhfr (ATCC Accession No. 37146). The plasmid contains the mouse DHFR gene under control of the SV40 early promoter. Chinese hamster ovary- or other cells lacking dihydrofolate activity that are transfected with these plasmids can be selected by growing the cells in a selective medium (alpha minus MEM, Life Technologies) supplemented with the

25 chemotherapeutic agent methotrexate. The amplification of the DHFR genes in cells resistant to methotrexate (MTX) has been well documented (see, e.g., Alt, F. W., Kellems, R. M., Bertino, J. R., and Schimke, R. T., 1978, *J Biol. Chem.* 253:1357-1370, Hamlin, J. L. and Ma, C. 1990, *Biochem. et Biophys. Acta*,

1097:107-143, Page, M. J. and Sydenham, M.A. 1991, *Biotechnology* 9:64-68).

Cells grown in increasing concentrations of MTX develop resistance to the drug by overproducing the target enzyme, DHFR, as a result of amplification of the DHFR gene. If a second gene is linked to the DHFR gene, it is usually co-amplified and over-expressed. It is known in the art that this approach may be used to develop cell lines carrying more than 1,000 copies of the amplified gene(s). Subsequently, when the methotrexate is withdrawn, cell lines are obtained which contain the amplified gene integrated into one or more chromosome(s) of the host cell.

Plasmid pC4 contains for expressing the gene of interest the strong promoter of the long terminal repeat (LTR) of the Rous Sarcoma Virus (Cullen, *et al.*, *Molecular and Cellular Biology*, March 1985:438-447) plus a fragment isolated from the enhancer of the immediate early gene of human cytomegalovirus (CMV) (Boshart *et al.*, *Cell* 41:521-530 (1985)). Downstream of the promoter are BamHI, XbaI, and Asp718 restriction enzyme cleavage sites that allow integration of the genes. Behind these cloning sites the plasmid contains the 3' intron and polyadenylation site of the rat preproinsulin gene. Other high efficiency promoters can also be used for the expression, e.g., the human β -actin promoter, the SV40 early or late promoters or the long terminal repeats from other retroviruses, e.g., HIV and HTLVI. Clontech's Tet-Off and Tet-On gene expression systems and similar systems can be used to express the BCSG1 in a regulated way in mammalian cells (Gossen, M., & Bujard, H. 1992, *Proc. Natl. Acad. Sci. USA* 89: 5547-5551). For the polyadenylation of the mRNA other signals, e.g., from the human growth hormone or globin genes can be used as well. Stable cell lines carrying a gene of interest integrated into the chromosomes can also be selected upon co-transfection with a selectable marker such as gpt, G418 or hygromycin. It is advantageous to use more than one selectable marker in the beginning, e.g., G418 plus methotrexate.

-42-

The plasmid pC4 is digested with the restriction enzymes BamHI/Asp718 and then dephosphorylated using calf intestinal phosphatase by procedures known in the art. The vector is then isolated from a 1% agarose gel.

5 The DNA sequence encoding the BCSG1 protein sequence is amplified using PCR oligonucleotide primers corresponding to the 5' and 3' sequences of the gene. The 5' primer has the sequence 5' GGGGATccgccaccATGTTTCAAGAAGG 3' (SEQ ID NO:7) (Kozak sequence is represented by the lowercase letters) containing the underlined BamHI restriction enzyme site followed by an efficient signal for initiation of translation in eukaryotes, as described by Kozak, M., *J. Mol. Biol.* 196:947-950 (1987), and 10 15 bases of the coding sequence of BCSG1 shown in Figure 1 (SEQ ID NO:1). The 3' primer has the sequence 5' GGGGTACCTCACTAGTCTCCCCCACTCTGG 3' (SEQ ID NO:9) containing the underlined Asp718 restriction site followed by 22 nucleotides complementary to the non-translated region of the BCSG1 gene shown in Figure 1 (SEQ ID 15 NO:1).

The amplified fragment is digested with the endonucleases BamHI/Asp718 and then purified again on a 1% agarose gel. The isolated fragment and the dephosphorylated vector are then ligated with T4 DNA ligase. 20 *E. coli* HB101 or XL-1 Blue cells are then transformed and bacteria are identified that contain the fragment inserted into plasmid pC4 using, for instance, restriction enzyme analysis.

Chinese hamster ovary cells lacking an active DHFR gene are used for transfection. 5 µg of the expression plasmid pC4 is cotransfected with 0.5 µg of 25 the plasmid pSV2-neo using lipofectin (Felgner *et al.*, *supra*). The plasmid pSV2neo contains a dominant selectable marker, the neo gene from Tn5 encoding an enzyme that confers resistance to a group of antibiotics including G418. The cells are seeded in alpha minus MEM supplemented with 1 mg/ml G418. After 2 days, the cells are trypsinized and seeded in hybridoma cloning plates (Greiner, 30 Germany) in alpha minus MEM supplemented with 10, 25, or 50 ng/ml of

metothrexate plus 1 mg/ml G418. After about 10-14 days single clones are trypsinized and then seeded in 6-well petri dishes or 10 ml flasks using different concentrations of methotrexate (50 nM, 100 nM, 200 nM, 400 nM, 800 nM). Clones growing at the highest concentrations of methotrexate are then transferred to new 6-well plates containing even higher concentrations of methotrexate (1 μ M, 2 μ M, 5 μ M, 10 mM, 20 mM). The same procedure is repeated until clones are obtained which grow at a concentration of 100 - 200 μ M. Expression of the desired gene product is analyzed, for instance, by SDS-PAGE and Western blot or by reverse phase HPLC analysis.

Example 4: Tissue distribution of BCSG1 mRNA expression

Northern blot analysis was carried out to examine BCSG1 gene expression in human tissues as follows. Total RNA was extracted from tissues according to the method of Chomczynski and Sacchi (Chomczynski, P. & Sacchi, N., *Anal. Biochem.* 162:156-159 (1987)). Equal aliquots of RNA were electrophoresed in a 1.2% agarose gel containing formaldehyde and transferred to nylon membrane (Boehringer Mannheim). The membrane was pre-hybridized with ExpressHyb hybridization solution (Clontech, Inc.) at 68°C for 30 min. The hybridization was carried out in the same solution with 32 P-labeled BCSG1 probe (1.5×10^6 cpm/ml) for 1 hour at 68°C. The membrane was then rinsed in 2 x SSC containing 0.05% SDS three times for 30 min at room temperature, followed by two washes with 0.1 x SSC containing 0.1% SDS for 40 min at 50°C. The full-length BCSG1 cDNA (SEQ ID NO:1) was isolated from the Bluescript vector, following EcoRI and XhoI digestion, and used as a template for preparation of a random-labelled cDNA probe. Random primer DNA labeling kit was obtained from Boehringer Mannheim, Indianapolis. 32 P-dATP was purchased from Amersham.

The northern blot showed that BCSG1 was abundantly expressed as the expected 1 kb transcript in brain which is a rich source for AD amyloid family

genes. A much less intense band of similar size was also seen in the following tissues: ovary, testis, colon, and heart.

Example 5: Cloning of BCSG1 from cDNA libraries

EST analysis was used to search for new genes differentially expressed in breast cancer versus normal breast. A data base containing approximately 500,000 human partial cDNA sequences (expressed sequence tags) has been established in a collaborative effort between the Institute for Genomic Research and Human Genome Science Inc., using high throughput automated DNA sequence analysis of randomly selected human cDNA clones (Adams, M.D., *et al.*, *Science* 252:1651-6 (1991)). RNAs from a stage III breast carcinoma and patient-matched normal breast were isolated and subjected to preparation of cDNA libraries. EST automated DNA sequence analysis was performed on randomly selected cDNA clones. Both libraries had about 60% novel gene sequences which did not match exactly to published human genes. A total of 3048 ESTs from breast cancer cDNA library and 2886 ESTs from normal breast cDNA library were randomly picked and sequence analyzed. The ESTs with overlapping sequences were grouped into unique EST groups; and each EST group may represent a gene or a family of sequence-related genes. There were more than 2,200 EST groups that were analyzed for quantitative comparison of EST hits in the pair of cDNA libraries from normal breast versus breast cancer by examining the expression of individual EST sequences. The numbers of EST hits in the libraries reflect the relative expression or mRNA transcript copy numbers of the EST. This direct differential cDNA sequence, as illustrated in Figure 2, utilizing the direct EST sequencing analysis simultaneously on a pair of cDNA libraries made from normal breast and breast cancer, was used to study expression profile of individual genes and patterns of genes in normal breast versus breast cancer.

Results

cDNA libraries were generated from breast cancer biopsy specimen and patient-matched normal breast and were analyzed by EST sequencing. Approximately 6,000 ESTs were analyzed and grouped to different groups based on sequence overlapping, and 2,200 unique EST groups were first analyzed for relative expression in the cDNA libraries from normal breast versus breast cancer and then subjected to tissue-specific expression by examining tissue origins of individual EST sequences against a large population of ESTs derived from a variety of different tissue types. Three classes of EST groups were identified that were differentially expressed in normal breast versus breast cancer. As a demonstration of this approach, Table 1 shows a partial list of three classes of genes that are differentially expressed in normal breast versus breast cancer. Class I represents the genes more abundant in breast cancer than in normal breast and includes cathepsin D, a well-studied steroid regulated extracellular matrix-degrading proteinase (Rocheft, H., *et al.*, *J. Cell. Biochem.* 35:17-29 (1987); Cavailles, V., *et al.*, *Biochem. Biophys. Res. Commun.* 174:816-24 (1991); Capony, F. *et al.*, *Biochem. Biophys. Res. Commun.* 171:972-80 (1990)). Cathepsin D is thought to play a role in breast cancer metastasis (Rocheft, H., *et al.*, *J. Cell. Biochem.* 35:17-29 (1987); Cavailles, V., *et al.*, *Biochem. Biophys. Res. Commun.* 174:816-24 (1991); Capony, F. *et al.*, *Biochem. Biophys. Res. Commun.* 171:972-80 (1990)) and has been proposed as a prognostic marker in breast cancer progression (Brouillet, J.P., *et al.*, *Eur. J. Cancer* 26:437-41 (1990); Spyrtos, F., *et al.*, *Lancet*, 11:1115-8 (1989); Rocheft, H., *et al.*, *J. steroid Biochem.* 34:177-82 (1989); Foekens J.A., *et al.*, *JNCI* 87:751-6 (1995)). As listed, there were 5 cathepsin D ESTs sequenced in the breast cancer cDNA library and only 1 EST in the normal breast cDNA library. Another proposed breast cancer metastasis-related gene and a prognostic marker for breast cancer, 67 kDa laminin receptor (Horan-Hand, P., *et al.*, *Cancer Res.* 45:833-40 (1986); Hunt, G. *Exp. Cell Biol* 57(3):165-76 (1989); Castronovo, V., *et al.*, *Am. J.*

Pathol. 137(6):1373-81 (1990); Marques, L.A., *et al.*, *Cancer Res.* 50(5):1479-83 (1990); Gasparini, G., *et al.*, *Int. J. Cancer.* 60(5):604-10 (1995)), was also picked up in this class by the Differential cDNA Sequencing approach. Class II represents genes that are more abundant in normal breast than in breast cancer.

5 Although the genes in classes I and II are differentially expressed in normal breast versus breast cancer, none of these genes are unique to breast tissues. Class III is a special group of genes that are selectively expressed in breast relative to other tissue types. The tissue-specific expression of the unique gene was searched against approximately 500,000 ESTs using the BLAST
10 program (Altschul, S.F., *et al.*, *J. Mol. Biol.* 215(3):403-10 (1990)). None of these breast cancer specific genes (BCSG) except the first one matched with any sequences in public gene sequence databases. BCSG1 was chosen for analysis as a first putative breast cancer maker gene because 1) its sequence has been matched with the sequence in public gene sequence database; and 2) most of the
15 individual EST sequences in BCSG1 derived from a breast tumor cDNA library. Of the eight distinctive EST clones in BCSG1, seven of them were discovered in breast cDNA libraries and only one in a brain library. Of the seven EST clones discovered in the breast cDNA libraries, six of them were identified in the breast tumor library and only one in the normal breast library. After complete
20 sequencing of all 6 EST clones, one EST clone was found to have a complete full-length sequence. The open reading frame of the resulting full-length gene is predicted to encode a 127 amino acid polypeptide. After optimal alignment, the putative BCSG1-encoded protein shows 54% sequence identity with the recently cloned non-A β fragment of human Alzheimer's disease (AD) amyloid protein
25 (Ueda, K., *et al.*, *Proc. Natl. Acad. Sci. USA.* 90(23):11282-6 (1993)).

-47-

**TABLE 1. PARTIAL LIST OF DIFFERENTIAL EXPRESSED GENES IN NORMAL VERSUS CANCEROUS
BREAST IDENTIFIED BY DIFFERENTIAL CDNA SEQUENCING**

Genes more abundant in breast cancer

5	Genes	EST	
		Cancer	Normal
	Breast basic conserved gene	33	9
	Cathepsin D	5	1
	67kDa laminin Receptor	4	0
10	Elongation factor 1	13	5

Genes More Abundant in Normal Breast

15	Genes	EST	
		Cancer	Normal
	Matrix Gla protein	0	8
	23 kDa Highly basic Protein	3	11

Genes as Breast-Specific and Differentially Expressed

20	Genes	EST	
		NB ¹	BC ²
	BCSG1	1	6
	BCSG2	0	7
	BCSG3	0	5
	BCSG4	4	0
	BCSG5	0	4

25 ¹ normal breast; ² breast cancer

30 Table 1. Complementary DNA libraries were established from a stage III breast carcinoma and patient-matched normal breast. A total of 5,934 ESTs were randomly picked and sequence analyzed. More than 2,200 distinctive EST groups were analyzed for quantitative comparison of EST hits in the pair of cDNA libraries from breast cancer versus normal breast as described in "Materials and Methods". The same EST groups were also analyzed by examining the tissue-specific expression against the total of 500,000 ESTs from a variety of different cDNA libraries. Only a unique EST group with more than 3 breast-specific EST hits was listed and the rest of the several dozens EST groups with fewer than 4 breast-specific EST hits were omitted in this list.

Example 6: Expression of BCSG1 in human breast cancer cells

In an attempt to evaluate the potential biological significance of BCSG1 on human breast cancer development and progression, BCSG1 gene expression in human breast biopsy samples was examined using Northern blot analysis.

5 The RNA from human breast cancer cells was prepared using the RNA isolation kit RNazol B (Tel-Test, Inc) based on the manufacturer's instruction. Equal aliquots of RNA were electrophoresed in a 1.2% agarose gel containing formaldehyde and transferred to nylon membrane (Boehringer Mannheim). The membrane was pre-hybridized with ExpressHyb hybridization solution (Clontech, Inc.) at 68°C for 30 min. The hybridization was carried out in the same solution with ³²P-labeled BCSG1 probe (1.5 x 10⁶ cpm/ml) for 1 hour at 68°C. The membrane was then rinsed in 2 x SSC containing 0.05% SDS three times for 30 min at room temperature, followed by two washes with 0.1 x SSC containing 0.1% SDS for 40 min at 50°C. The full-length BCSG1 cDNA (SEQ ID NO:1) was isolated from the Bluescript vector, following EcoRI and XhoI digestion, and used as a template for preparation of a random-labelled cDNA probe. Random primer DNA labeling kit was obtained from Boehringer Mannheim, Indianapolis. ³²P-dATP was purchased from Amersham.

15 The expression of BCSG1 in metastatic breast carcinoma and benign breast tissue were analyzed by Northern blotting. Overexpression of the BCSG1 transcript in breast carcinoma. In contrast, the BCSG1 transcript was undetectable in benign breast tissue. The presence of BCSG1 transcript in human breast tissue and its overexpression in breast carcinomas are consistent with the differential cDNA sequencing cloning strategy which suggests a possible role or a biomarker of up-regulation of BCSG1 in the development of breast cancer.

20 The expression of BCSG1 was also examined in a variety of human breast cancer cell lines, namely, primary solid tumor derived cell lines H3477, H3630, H3680B; pleural effusion derived cell lines H3396, MCF7, SKBR-3 MDAMB231; infiltrating ductal carcinoma derived cell lines H3914, H3922, ZR-

75-1, T47D. Cell lines of T47D, ZR-75-1, SKBR-3, MCF-7 and MDA-MB-231 are from ATCC; all other lines were initially isolated at Bristol-Myers Squibb Pharmaceutical Research Institute (Liu, J., *Cancer Res.*).

5 Northern blot detected the 1 Kb BCSG1 transcript in 2 of the 4 cell lines derived from pleural effusion (i.e., SKBR-3 MDAMB231) and all 4 cell lines detected from ductal infiltrating carcinomas. Interestingly, none of the cell lines derived from primary solid breast carcinoma expressed BCSG1 mRNA. Among these lines, H3922 expressed the highest level of BCSG1 mRNA. The absence of BCSG1 mRNA in some pleural effusion derived cell lines suggest that the
10 expression of BCSG1 gene may require specific *in vivo* conditions, or that it is induced by interactions between the tumor cells and stromal cells.

Example 7: In situ hybridization of BCSG1 in breast cancer cells

In order to localize the cellular source of the BCSG1 expression and to further assess the biological relevance of the overexpression of BCSG1 in breast
15 cancers, *in situ* hybridization was performed on fixed breast sections from 20 infiltrating carcinomas, 15 *in situ* carcinomas, and 15 benign breast lesions (breast hyperplasia and fibroadenoma).

In situ hybridization was carried out as described (M.L. Angerer & R.C. Angerer, In: *In situ hybridization*, D. Rickwood and B.D. Hames (ed.). London: LRL Press., (1992), pp. 15-32). Briefly, deparaffinized and acid-treated sections
20 (5-um thick) were treated with proteinase K, pre-hybridized, and hybridized overnight with digoxigenin labeled anti-sense transcripts from a BCSG1 cDNA insert. The BCSG1 antisense probe is a 550 bp full-length fragment. The probe was generated by PstI cut of BCSG1 cDNA plasmid and followed by T7
25 polymerase. Hybridization was followed by RNase treatment and three stringent washings. Sections were incubated with mouse anti-digoxigenin antibodies (Boehringer) followed by the incubation with biotin-conjugated secondary rabbit anti-mouse antibodies (DAKO). The calorimetric detection were performed

-50-

using a standard indirect streptavidin-biotin immunoreaction method by DAKO's Universal LSAB Kit according to manufacturer's instructions.

In these experiments, two aspects of BCSG1 expression were examined:

1) the tissue localization (stromal versus epithelial); and 2) the correlation of

5 BCSG1 expression and breast cancer malignant phenotype. A strongly positive BCSG1 hybridization in neoplastic epithelial cells of highly infiltrating breast carcinomas was observed. The expression of BCSG1 mRNA was detectable in the neoplastic epithelial cells in 18 of 20 infiltrating breast carcinomas. No expression of BCSG1 was detected in the stromal cells. In contrast, expression
10 of BCSG1 was absent in all 15 cases of normal or benign breast lesions. A representative negative staining of BCSG1 in an atypical proliferative breast lesion, a benign fibroadenoma, and normal ductal breast epithelial cells are presented. Furthermore, in a highly invasive breast carcinoma, no detectable signal of BCSG1 expression was evident in the residual normal lobular breast
15 epithelial cells although the surrounding invasive breast carcinoma cells were stained positive for BCSG1 expression. These *in situ* hybridization results are consistent with the Northern blot analysis which showed a strong expression of BCSG1 transcript in breast carcinoma but not in normal or benign breast lesions.

It is interesting to note that although a strong BCSG1 signal was easily
20 detected in the malignant breast epithelial cells of infiltrating breast carcinoma, the low grade *in situ* carcinomas showed a sparse and a light staining. Among 15 *in situ* carcinomas, 9 specimens were stained negatively and 6 specimens were partially stained. Both the intensity of staining and the number of positive cells were significantly reduced in the *in situ* breast carcinomas compared to the strong
25 expression in the metastatic breast carcinomas. These results, which demonstrated a gradient and stage-specific BCSG1 expression from virtually no detectable expression in normal or benign breast to the low level and partial expression in the low grade *in situ* breast carcinoma and to the high expression in the infiltrating malignant breast carcinomas, suggest an association of BCSG1

-51-

expression with breast cancer malignant progression. Based on this BCSG1 expression pattern, BCSG1 is useful as a breast cancer progression marker.

It will be clear that the invention may be practiced otherwise than as particularly described in the foregoing description and examples.

5 Numerous modifications and variations of the present invention are possible in light of the above teachings and, therefore, are within the scope of the appended claims.

10 The entire disclosure of all publications (including patents, patent applications, journal articles, laboratory manuals, books, or other documents) cited herein are hereby incorporated by reference.

-52-

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: HUMAN GENOME SCIENCES, INC.
9410 KEY WEST AVENUE
ROCKVILLE, MD 20850
UNITED STATES OF AMERICA

APPLICANT/INVENTOR: Ji, Hongjun
13144 Country Ridge Dr.
Germantown, MD 20874
United States of America

Rosen, Craig A.
22400 Rolling Hill Road
Laytonsville, MD 20882
United States of America

(ii) TITLE OF INVENTION: Breast Cancer Specific Gene 1

(iii) NUMBER OF SEQUENCES: 11

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Sterne, Kessler, Goldstein & Fox P.L.L.C.
(B) STREET: 1100 New York Ave., Suite 600
(C) CITY: Washington
(D) STATE: DC
(E) COUNTRY: USA
(F) ZIP: 20005-3934

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER: TO BE ASSIGNED
(B) FILING DATE: HEREWITH
(C) CLASSIFICATION:

(vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: US 60/037,080
(B) FILING DATE: 03-FEB-1997
(C) CLASSIFICATION:

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Steffe, Eric K.
(B) REGISTRATION NUMBER: 36,688
(C) REFERENCE/DOCKET NUMBER: 1488.081PC03

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: 202-371-2600
(B) TELEFAX: 202-371-2540

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 550 base pairs
(B) TYPE: nucleic acid

-53-

(C) STRANDEDNESS: double
(D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

(A) NAME/KEY: CDS
(B) LOCATION: 12..392

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

CACGAGCCAC C	ATG GAT GTT TTC AAG AAG GGC TTC TCC ATC GCC AAG AAG	50
	Met Asp Val Phe Lys Lys Gly Phe Ser Ile Ala Lys Lys	
	1 5 10	
GGC GTG GTG GGT GCG GTG GAA AAG ACC AAG CAG GGG GTG ACG GAA GCA		98
Gly Val Val Gly Ala Val Glu Lys Thr Lys Gln Gly Val Thr Glu Ala		
	15 20 25	
GCT GAG AAG ACC AAG GAG GGG GTC ATG TAT GTG GGA GCC AAG ACC AAG		146
Ala Glu Lys Thr Lys Glu Gly Val Met Tyr Val Gly Ala Lys Thr Lys		
	30 35 40 45	
GAG AAT GTT GTA CAG AGC GTG ACC TCA GTG GCC GAG AAG ACC AAG GAG		194
Glu Asn Val Val Gln Ser Val Thr Ser Val Ala Glu Lys Thr Lys Glu		
	50 55 60	
CAG GCC AAC GCC GTG AGC AAG GCT GTG GTG AGC AGC GTC AAC ACT GTG		242
Gln Ala Asn Ala Val Ser Lys Ala Val Val Ser Ser Val Asn Thr Val		
	65 70 75	
GCC ACC AAG ACC GTG GAG GAG GCG GAG AAC ATC GCG GTC ACC TCC GGG		290
Ala Thr Lys Thr Val Glu Glu Ala Glu Asn Ile Ala Val Thr Ser Gly		
	80 85 90	
GTG GTG CGC AAG GAG GAC TTG AGG CCA TCT GCC CCC CAA CAG GAG GGT		338
Val Val Arg Lys Glu Asp Leu Arg Pro Ser Ala Pro Gln Gln Glu Gly		
	95 100 105	
GAG GCA TCC AAA GAG AAA GAG GAA GTG GCA GAG GAG GCC CAG AGT GGG		386
Glu Ala Ser Lys Glu Lys Glu Glu Val Ala Glu Glu Ala Gln Ser Gly		
	110 115 120 125	
GGA GAC TAGAGGGCTA CAGGCCAGCG TGGATGACCT GAAGAGCGCT CCTCTGCCTT		442
Gly Asp		
GGACACCATC CCCTCCTAGC ACAAGGAGTG CCCGCCTTGA GTGACATGCG GGTGCCCACG		502
CTCCTGCCCT CGTCTCCCTG GACACCCTTG GCCTGTCCAC CTGTGCTG		550

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 127 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

-54-

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Asp Val Phe Lys Lys Gly Phe Ser Ile Ala Lys Lys Gly Val Val
 1 5 10 15
 Gly Ala Val Glu Lys Thr Lys Gln Gly Val Thr Glu Ala Ala Glu Lys
 20 25 30
 Thr Lys Glu Gly Val Met Tyr Val Gly Ala Lys Thr Lys Glu Asn Val
 35 40 45
 Val Gln Ser Val Thr Ser Val Ala Glu Lys Thr Lys Glu Gln Ala Asn
 50 55 60
 Ala Val Ser Lys Ala Val Val Ser Ser Val Asn Thr Val Ala Thr Lys
 65 70 75 80
 Thr Val Glu Glu Ala Glu Asn Ile Ala Val Thr Ser Gly Val Val Arg
 85 90 95
 Lys Glu Asp Leu Arg Pro Ser Ala Pro Gln Gln Glu Gly Glu Ala Ser
 100 105 110
 Lys Glu Lys Glu Glu Val Ala Glu Glu Ala Gln Ser Gly Gly Asp
 115 120 125

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 23 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

GGGGATCCAT GTTTTCAAGA AGG

23

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 27 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

GGAAGCTTCT AGTCTCCCCC ACTCTGG

27

(2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:

-55-

- (A) LENGTH: 25 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

GGGGATCCCG ATGTTTTCAA GAAGG

25

(2) INFORMATION FOR SEQ ID NO:6:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 27 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

GGGGTACCCT AGTCTCCCC ACTCTGG

27

(2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 29 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

GGGGATCCGC CACCATGTTT TCAAGAAGG

29

(2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 60 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

-56-

G3GGATCCTC AGAAAGCGTA GTCTGGGACG TCGTATGGGT ACTAGTCTCC CCCACTCTGG 60

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 30 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

GGGGTACCTC ACTAGTCTCC CCCACTCTGG 30

(2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 3974 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: both
(D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

GGTACCTAAG TGAGTAGGGC GTCCGATCGA CGGACGCCCT TTTTTGAAT TCGTAATCAT 60
GGTCATAGCT GTTTCCTGTG TGAAATTGTT ATCCGCTCAC AATCCACAC AACATACGAG 120
CCGGAAGCAT AAAGTGTAAG GCCTGGGGTG CCTAATGAGT GAGCTAACTC ACATTAATTG 180
CGTTGCGCTC ACTGCCCCGCT TTCCAGTCGG GAAACCTGTC GTGCCAGCTG CATTAAATGAA 240
TCGGCCAACG CGCGGGGAGA GGCGGTTTGC GTATTGGGCG CTCTTCCGCT TCCTCGCTCA 300
CTGACTCGCT GCGCTCGGTC GTTCGGCTGC GGCGAGCGGT ATCAGCTCAC TCAAAGGCGG 360
TAATACGGTT ATCCACAGAA TCAGGGGATA ACGCAGGAAA GAACATGTGA GCAAAGGCC 420
AGCAAAGGC CAGGAACCGT AAAAAGGCCG CGTTGCTGGC GTTTTCCAT AGGCTCCGCC 480
CCCCTGACGA GCATCAGAAA AATCGACGCT CAAGTCAGAG GTGGCGAAAC CCGACAGGAC 540
TATAAGATA CCAGGCGTTT CCCCTGGAA GCTCCCTCGT GCGCTCTCCT GTTCCGACCC 600
TGCCGCTTAC CGGATACCTG TCCGCTTTC TCCCTTCGGG AAGCGTGGCG CTTTCTCATA 660
GCTCACGCTG TAGGTATCTC AGTTCGGTGT AGGTCGTTCT CTCCAAGCTG GGCTGTGTGC 720
ACGAACCCCC CGTTCAGCCC GACCGCTGCG CCTTATCCGG TAACTATCGT CTTGAGTCCA 780
ACCCGGTAAG ACACGACTTA TCGCCACTGG CAGCAGCCAC TGGTAACAGG ATTAGCAGAG 840

-57-

CGAGGTATGT AGGCGGTGCT ACAGAGTTCT TGAAGTGGTG GCCTAACTAC GGCTTACACTA	900
GAAGAACAGT ATTTGGTATC TGCCTCTGCT TGAAGCCAGT TACCTTCGGA AAAAGAGTTG	960
GTAGCTCTTG ATCCGGCAAA CAAACCACCG CTGGTAGCGG TGGTTTTTTT GTTTGCAAGC	1020
AGCAGATTAC GCGCAGAAAA AAAGGATCTC AAGAAGATCC TTTGATCTTT TCTACGGGGT	1080
CTGACGCTCA GTGGAACGAA AACTCACGTT AAGGGATTTT GGTTCATGAGA TTATCGTCGA	1140
CAATTCGCGC GCGAAGGCGA AGCGGCATGC ATTTACGTTG ACACCATCGA ATGGTGCAAA	1200
ACCTTTTCGCG GTATGGCATG ATAGCGCCCG GAAGAGAGTC AATTCAGGGT GGTGAATGTG	1260
AAACCAGTAA CGTTATACGA TGTCGCAGAG TATGCCGGTG TCTCTTATCA GACCGTTTCC	1320
CGCGTGGTGA ACCAGGCCAG CCACGTTTCT GCGAAAACGC GGGAAAAAGT GGAAGCGGCG	1380
ATGGCGGAGC TGAATTACAT TCCCAACCGC GTGGCACAAC AACTGGCGGG CAAACAGTCG	1440
TTGCTGATTG GCGTTGCCAC CTCCAGTCTG GCCCTGCACG CGCCGTCGCA AATTGTCGCG	1500
GCGATTAAAT CTCGCGCCGA TCAACTGGGT GCCAGCGTGG TGGTGTCGAT GGTAGAACGA	1560
AGCGGCGTCG AAGCCTGTAA AGCGGCGGTG CACAATCTTC TCGCGCAACG CGTCAGTGGG	1620
CTGATCATTG ACTATCCGCT GGATGACCAG GATGCCATTG CTGTGGAAGC TGCCTGCACT	1680
AATGTTCCGG CGTTATTTCT TGATGTCTCT GACCAGACAC CCATCAACAG TATTATTTTC	1740
TCCCATGAAG ACGGTACGCG ACTGGGCGTG GAGCATCTGG TCGCATTGGG TCACCAGCAA	1800
ATCGCGCTGT TAGCGGGCCC ATTAAGTTCT GTCTCGGCGC GTCTGCGTCT GGCTGGCTGG	1860
CATAAATATC TCACTCGCAA TCAAATTCAG CCGATAGCGG AACGGGAAGG CGACTGGAGT	1920
GCCATGTCCG GTTTTCAACA AACCATGCAA ATGCTGAATG AGGGCATCGT TCCCACTGCG	1980
ATGCTGGTTG CCAACGATCA GATGGCGCTG GGCGCAATGC GCGCCATTAC CGAGTCCGGG	2040
CTGCGCGTTG GTGCGGATAT CTCGGTAGTG GGATACGACG ATACCGAAGA CAGCTCATGT	2100
TATATCCCGC CGTTAACCAC CATCAAACAG GATTTTCGCC TGCTGGGGCA AACCAGCGTG	2160
GACCGCTTGC TGCAACTCTC TCAGGGCCAG GCGGTGAAGG GCAATCAGCT GTTGCCCGTC	2220
TCACTGGTGA AAAGAAAAAC CACCCTGGCG CCCAATACGC AAACCGCCTC TCCCCGCGCG	2280
TTGGCCGATT CATTAATGCA GCTGGCACGA CAGGTTTCCC GACTGGAAAG CGGGCAGTGA	2340
GCGCAACGCA ATTAATGTAA GTTAGCGCGA ATTGTCGACC AAAGCGGCCA TCGTGCCTCC	2400
CCACTCCTGC AGTTCGGGGG CATGGATGCG CGGATAGCCG CTGCTGGTTT CCTGGATGCC	2460
GACGGATTG CACTGCCGGT AGAACTCCGC GAGGTCGTCC AGCCTCAGGC AGCAGCTGAA	2520
CCAACTCGCG AGGGGATCGA GCCCGGGGTG GGCGAAGAAC TCCAGCATGA GATCCCCGCG	2580
CTGGAGGATC ATCCAGCCGG CGTCCCGGAA AACGATTCCG AAGCCCAACC TTTCATAGAA	2640
GGCGGCGGTG GAATCGAAAT CTCGTGATGG CAGGTTGGGC GTCGCTTGGT CGGTCAATTC	2700
GAACCCGAGA GTCCCGCTCA GAAGAACTCG TCAAGAAGGC GATAGAAGGC GATGCGCTGC	2760

-58-

GAATCGGGAG CGGCGATACC GTAAAGCAGC AGGAAGCGGT CAGCCCATTG GCCGCCAAGC	2820
TCTTCAGCAA TATCACGGGT AGCCAACGCT ATGTCCTGAT AGCGGTCCGC CACACCCAGC	2880
CGGCCACAGT CGATGAATCC AGAAAAGCGG CCATTTTCCA CCATGATATT CGGCAAGCAG	2940
GCATCGCCAT GGGTCACGAC GAGATCCTCG CCGTCGGGCA TGC GCGCCTT GAGCCTGGCG	3000
AACAGTTTCGG CTGGCGCGAG CCCCTGATGC TCTTCGTCCA GATCATCCTG ATCGACAAGA	3060
CCGGCTTCCA TCCGAGTAGC TGCTCGCTCG ATGCGATGTT TCGCTTGGTG GTCGAATGGG	3120
CAGGTAGCCG GATCAAGCGT ATGCAGCCGC CGCATTGCAT CAGCCATGAT GGATACTTTC	3180
TCGGCAGGAG CAAGGTGAGA TGACAGGAGA TCCTGCCCCG GCACTTCGCC CAATAGCAGC	3240
CAGTCCCTTC CCGCTTCAGT GACAACGTCG AGCACAGCTG CGCAAGGAAC GCCCGTCGTG	3300
GCCAGCCACG ATAGCCGCGC TGCCTCGTCC TGCACTTCAT TCAGGGCACC GGACAGGTCG	3360
GTCTTGACAA AAAGAACCGG GCGCCCTGC GCTGACAGCC GGAACACGGC GGCATCAGAG	3420
CAGCCGATTG TCTGTTGTGC CCAGTCATAG CCGAATAGCC TCTCCACCA AGCGGCCGGA	3480
GAACCTGCGT GCAATCCATC TTGTTCAATC ATGCGAAACG ATCCTCATCC TGTCTCTTGA	3540
TCAGATCTTG ATCCCCTGCG CCATCAGATC CTTGGCGGCA AGAAAGCCAT CCAGTTTACT	3600
TTGCAGGGCT TCCCAACCTT ACCAGAGGGC GCCCCAGCTG GCAATCCGG TTCGCTTGCT	3660
GTCCATAAAA CCGCCAGTC TAGCTATCGC CATGTAAGCC CACTGCAAGC TACCTGCTTT	3720
CTCTTTGCGC TTGCGTTTTC CCTTGTCAG ATAGCCAGT AGCTGACATT CATCCGGGGT	3780
CAGCACCGTT TCTGCGGACT GGCTTTCTAC GTGTTCCGCT TCCTTTAGCA GCCCTTGCGC	3840
CCTGAGTGCT TGCGGCAGCG TGAAGCTTAA AAACTGCAA AAAATAGTTT GACTGTGAG	3900
CGGATAACAA TTAAGATGTA CCCAATTGTG AGCGGATAAC AATTTCACAC ATTAAAGAGG	3960
AGAAATTACA TATG	3974

(2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 112 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: both
 - (D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

AAGCTTAAAA AACTGCAAAA AATAGTTTGA CTTGTGAGCG GATAACAATT AAGATGTACC	60
CAATTGTGAG CGGATAACAA TTTCACACAT TAAAGAGGAG AAATTACATA TG	112

- 59 -

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM
(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page ____3____, line ____19____.	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input checked="" type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 12301 Parklawn Drive Rockville, Maryland 20852 United States of America	
Date of deposit June 2, 1995	Accession Number 97175
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
DNA Plasmid, PF 158	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the international Bureau later (specify the general nature of the indications, e.g., "Accession Number of Deposit")	

For receiving Office use only	For International Bureau use only
<input type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer	Authorized officer

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM
(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page ____3____, line ____19____.	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 12301 Parklawn Drive Rockville, Maryland 20852 United States of America	
Date of deposit January 23, 1997	Accession Number 97856
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
DNA Plasmid BCSG	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the international Bureau later (specify the general nature of the indications, e.g., "Accession Number of Deposit")	

For receiving Office use only	For International Bureau use only
<input type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer	Authorized officer

- 61 -

What Is Claimed Is:

1. An isolated nucleic acid molecule comprising a polynucleotide having a nucleotide sequence at least 95% identical to a sequence selected from the group consisting of:
 - 5 (a) a nucleotide sequence encoding a polypeptide comprising amino acids from about 1 to about 127 in SEQ ID NO:2;
 - (b) a nucleotide sequence encoding a polypeptide comprising amino acids from about 2 to about 127 in SEQ ID NO:2;
 - (c) a nucleotide sequence encoding a polypeptide having the
10 amino acid sequence encoded by the cDNA clones contained in ATCC Deposit No. 97175 or 97856; and
 - (d) a nucleotide sequence complementary to any of the nucleotide sequences in (a), (b) or (c).
2. An isolated nucleic acid molecule comprising a polynucleotide
15 which encodes the amino acid sequence of an epitope-bearing portion of an BCSG1 polypeptide having an amino acid sequence in (a), (b), (c) or (d) of claim 1.
3. The isolated nucleic acid molecule of claim 7, which encodes an epitope-bearing portion of a BCSG1 polypeptide selected from the group
20 consisting of: a polypeptide comprising amino acid residues from about 94 to about 107 in Figure 1 (SEQ ID NO:2); and a polypeptide comprising amino acid residues from about 120 to about 127 in Figure 1 (SEQ ID NO:2).
4. An isolated nucleic acid molecule, comprising a polynucleotide
25 having a sequence selected from the group consisting of:

- 62 -

(a) the nucleotide sequence of a fragment of the sequence shown in SEQ ID NO:1, wherein said fragment comprises at least 50 contiguous nucleotides of SEQ ID NO:1; and

5 (b) a nucleotide sequence complementary to a nucleotide sequence in (a).

5. A method for making a recombinant vector comprising inserting an isolated nucleic acid molecule of claim 1 into a vector.

6. A recombinant vector produced by the method of claim 5.

10 7. A method of making a recombinant host cell comprising introducing the recombinant vector of claim 6 into a host cell.

8. A recombinant host cell produced by the method of claim 7.

15 9. A recombinant method for producing any of the BCSG1 polypeptides, comprising culturing the recombinant host cell of claim 8 under conditions such that said polypeptide is expressed and recovering said polypeptide.

10. An isolated BCSG1 polypeptide having an amino acid sequence at least 95% identical to a sequence selected from the group consisting of:

- 20 (a) amino acids from about 1 to about 127 in SEQ ID NO:2;
(b) amino acids from about 2 to about 127 in SEQ ID NO:2;
(c) the amino acid sequence of the BCSG1 polypeptide having the amino acid sequence encoded by the cDNA clones contained in ATCC Deposit No. 97856 or 97175; and
25 (d) the amino acid sequence of an epitope-bearing portion of any one of the polypeptides of (a), (b) or (c).

- 63 -

11. An isolated polypeptide comprising an epitope-bearing portion of the BCSG1 protein, wherein said portion is selected from the group consisting of: a polypeptide comprising amino acid residues from about 94 to about 107 in Figure 1 (SEQ ID NO:2); and a polypeptide comprising amino acid residues from about 120 to about 127 in Figure 1 (SEQ ID NO:2).

12. An isolated antibody that binds specifically to a BCSG1 polypeptide of claim 10.

13. An isolated nucleic acid molecule comprising a polynucleotide encoding a BCSG1 polypeptide wherein, except for one to fifty conservative amino acid substitutions, said polypeptide has a sequence selected from the group consisting of:

(a) a nucleotide sequence encoding a polypeptide comprising amino acids from about 1 to about 127 in SEQ ID NO:2;

(b) a nucleotide sequence encoding a polypeptide comprising amino acids from about 2 to about 127 in SEQ ID NO:2;

(c) a nucleotide sequence encoding a polypeptide having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 97175 or 97856; and

(d) a nucleotide sequence complementary to any of the nucleotide sequences in (a), (b) or (c).

14. An isolated BCSG1 polypeptide wherein, except for at least one conservative amino acid substitution, said polypeptide has a sequence selected from the group consisting of:

(a) amino acids from about 1 to about 127 in SEQ ID NO:2;

(b) amino acids from about 2 to about 127 in SEQ ID NO:2;

- 64 -

(c) the amino acid sequence of the BCSG1 polypeptide having the amino acid sequence encoded by the cDNA clone contained in ATCC Deposit No. 97175 or 97856; and

5 (d) the amino acid sequence of an epitope-bearing portion of any one of the polypeptides of (a), (b) or (c).

15. A method for breast tumor diagnosis in an individual comprising assaying the expression level of the gene encoding the BCSG1 protein in cells or body fluid of the individual and comparing the gene expression level with a standard BCSG1 gene expression level, whereby an increase in the gene
10 expression level over the standard is indicative of malignant breast cancer.

1/4

10 30 50
CACGAGCCACCATGGATGTTTTCAAGAAGGGCTTCTCCATCGCCAAGAAGGGCGTGGTGG
M D V F K K G F S I A K K G V V G

70 90 110
GTGCGGTGGAAGACCAAGCAGGGGGTGACGGAAGCAGCTGAGAAGACCAAGGAGGGGG
A V E K T K Q G V T E A A E K T K E G V

130 150 170
TCATGTATGTGGGAGCCAAGACCAAGGAGAATGTTGTACAGAGCGTGACCTCAGTGGCCG
M Y V G A K T K E N V V Q S V T S V A E

190 210 230
AGAAGACCAAGGAGCAGGCCAACGCCGTGAGCAAGGCTGTGGTGAGCAGCGTCAACACTG
K T K E Q A N A V S K A V V S S V N T V

250 270 290
TGGCCACCAAGACCGTGGAGGAGGCGGAGAACATCGCGGTACCTCCGGGGTGGTGCACA
A T K T V E E A E N I A V T S G V V R K

310 330 350
AGGAGGACTTGAGGCCATCTGCCCCCAACAGGAGGGTGAGGCATCCAAAGAGAAAGAGG
E D L R P S A P Q Q E G E A S K E K E E

370 390 410
AAGTGGCAGAGGAGGCCAGAGTGGGGGAGACTAGAGGGCTACAGGCCAGCGTGGATGAC
V A E E A Q S G G D *

430 450 470
CTGAAGAGCGCTCCTCTGCCTTGACACCATCCCCTCCTAGCACAAAGAGTGCCCGCCTT

490 510 530
GAGTGACATGCGGGTGCCACGCTCCTGCCCTCGTCTCCCTGGACACCCTTGGCCTGTCC

550
ACCTGTGCTG

FIG. 1

2/4

DIFFERENTIAL cDNA SEQUENCING




FIG.2




FIG.3

4/4

1 AACCTT AAAAACTGCAAAAAATAGTTTGACTTTCGACCGGATAACAAT -35 OPERATOR 1

50 TAAGATGTACCCAATTGTGAGCGGATAACAAATTCACACATTAA -10 OPERATOR 2

94 S/D
ACAGCAGAAATTATCATATC

FIG.4

INTERNATIONAL SEARCH REPORT

Int. Application No
PCT/US 98/01804

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/12 C07K14/47 C07K16/18 C12N15/63 C12N5/10
C12N1/21 C12Q1/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C07K C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 97 02280 A (HUMAN GENOME SCIENCES, INC.) 23 January 1997 see page 1, paragraph 1 see page 2, paragraph 2 - page 3, paragraph 3 see page 6, paragraph 2 - paragraph 4 see page 7, paragraph 2 - page 10, paragraph 4 see page 12, last paragraph - page 59, paragraph 1; figure 1	1-15
A	MARK A. WATSON ET AL.: "Isolation of differentially expressed sequence tags from human breast cancer" CANCER RESEARCH, vol. 54, no. 17, 1 September 1994, MD US, pages 4598-4602, XP000576043 see the whole document	1-15
-/--		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

19 May 1998

Date of mailing of the international search report

03/06/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo rd.
Fax: (+31-70) 340-3016

Authorized officer

Montero Lopez, B

INTERNATIONAL SEARCH REPORT

Int. l. Application No

PCT/US 98/01804

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	KENJI UÉDA ET AL.: "Molecular cloning of cDNA encoding an unrecognized component of amyloid in Alzheimer disease" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, vol. 90, December 1993, WASHINGTON US, pages 11282-11286, XP002065401 cited in the application -----	
A	SHI Y E ET AL: "Identification of a putative breast cancer progression marker, BCM1, by differential cDNA sequencing." 87TH ANNUAL MEETING OF THE AMERICAN ASSOCIATION FOR CANCER RESEARCH, WASHINGTON, D.C., USA, APRIL 20-24, 1996. PROCEEDINGS OF THE AMERICAN ASSOCIATION FOR CANCER RESEARCH ANNUAL MEETING 37 (0). 1996. 238. ISSN: 0197-016X, XP002065402 see abstract no. 1628 -----	1-15
P,X	HONGJUN JI ET AL.: "Identification of a breast cancer-specific gene, BCSG1, by direct differential cDNA sequencing" CANCER RESEARCH, vol. 57, no. 4, 15 February 1997, MD US, pages 759-764, XP002065403 see the whole document -----	1-15

Information on patent family members

PCT/US 98/01804

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9702280 A	23-01-1997	AU 3092995 A	05-02-1997